

### Additional information

---

#### Physical features of the site:

##### *Geology, hydrogeology*

Following the withdrawal and the sedimentation of the last lake, the so-called Late Miocene Lake Pannon in this region on the Great Plain, approximately 4.5 million years ago, the ancestor structures of the Danube, the ancient Tisza and the tributaries of the latter appeared. From this point on the previous lake sediment supply was replaced by river sedimentation (primarily by the Danube). Until the Günz-Minden Interglacial Episode in the Pleistocene Ice Age following Pliocene Epoch the Danube ran southeast-bound towards Szeged, cutting the region in half, and supplied river sedimentation in a width of some 1000 metres. In the Günz-Minden Interglacial Episode of the Ice Age a major change occurred: with the development of the region's south-western depression (Kalocsa depression) the Danube gradually started to drift westward by leaving its previous diagonal flow direction and took over its present north-south position. The Danube had already filled up the previous areas. River sedimentation ceased on the alluvial fan replacing these, situated east of the region, which remained higher than the Trans-Tisza region, and a thick eolic sedimentary layer were deposited on it (in the areas undisturbed by water).

This sedimentary layer consists of sand blown out of the Danube valley in the ice-free periods of the Ice Age, which was structured as a series of sand piles in the north-south direction according to the dominant wind direction, as well as loess developed during the ice formation periods, their transformed (e.g. soil) varieties and sediments washed out by local precipitation.

The sediment pattern delivered by the Danube-Tisza interfluvial winds protrudes slightly east of the current Tisza route, between the river layers of the Tisza. Therefore a geological situation developed in the smaller eastern section of the region where the Tisza, through its westbound movement, entered the alluvial fan of Danubian origin and in certain locations cut up and destroyed the surface of Danubian origin from the late Pleistocene period and enriched it with its own sediments (occasionally in an astonishing width of several hundred metres).

Based on geological evolution, the geological structures covering the surface and the morphological conditions the region can be divided into three major geological units:

- Danube Valley (a tectonic and erosional depression along the Danube river in a width of some 20-30 km) with an average height of 90-100 m above sea level.

##### *Pedology*

Prior to the river control of the Danube the Danube Valley used to be the river's normal floodplain, then it was an area covered with inland waters on a regular basis subsequently, as well. Also, as a result of its pedological (mainly calcareous-sodic plains developed on a fine granule rock bed) and geological structure (the significant presence of a fine waterproof clay layer) precipitation filter downwards with difficulty and may remain permanently in the depressions. It is generally true that due to the winter precipitation and the high ground water level in the spring significant water volumes appear in the depressed areas (in the isolated depressions of lake beds and old water flows).

The soil types developed here are:

- Chernozem meadow soil types, which are surfaces developed on a sandy loess base situated in the highest level layers in the region, with a high humus content. Their layer thickness varies between 20-40 cm. Generally the salty ground water already does not impregnate these layers. In cases where these highest locations are relatively expansive, tillage activities are carried out on them, and if they are smaller in size (a few 100 m<sup>2</sup>), they form islands on the saline steppe, partly conserving the old sand and loess steppe flora of these areas.
- Solonetz meadow or carbonated solonetz soils, which appear in non-classical forms, in patches, and are more of a transition between the meadow and sodic soils in various combinations,
- Solonchak-solonetz soils, sodic solonchak soils, solonchak soils of eroded salt berms. Among these calcareous-sodic solonchak-solonetz soils are the most common, giving the character of the sodic plains found here. Due to their isolation and weak runoff conditions such depressions and low areas promoted the accumulation of periodic waters, which, as a result of the known salt composition of ground water, led to the formation of sodic wetlands and higher level sodic areas.

### *Climate*

The climate variations are limited in the region of the Carpathian Basin. The macroclimate can be considered a homogenous basic feature in terms of surface and fauna evolution, as well.

The region has a temperate continental climate. Its unique features are limited cloudiness, a relatively high number of sunshine hours, high daily and annual temperature variation, relative dryness and very low humidity values.

This region is the area with the least cloudiness in Hungary. The annual average cloudiness is 52-57%. The annual average number of sunshine hours is approx. 2050. At the same time this is one of the warmest areas in the country. No significant variations exist in this region. The annual average temperature is between 10-11°C. The mean temperature of the coldest month (January) is between minus 1.5 and minus 2°C, while that of the warmest month (July) is 21-22°C. Characteristically of areas with a continental climate, the annual average temperature variance is quite significant (23-24°C). The region can be classified within Hungary as one with a short winter and a long summer. The number of winter days is only 26-31, however, major frosts are common. Spring comes early, and the average temperature rises above 10°C in the whole region between 7-12 March. The number of summer days is 81-84. In the fall the daily average temperature falls below 10°C again generally between 17-21 October.

The region is one of the parts of the country having the least precipitation. Under normal conditions the annual precipitation is between 500-600 mm in the region. The rainfall of the summer semester (April-September), the so-called breeding period, is around 300-350 mm. The winter precipitation occurs mainly in the form of snow. The number of snow-cover days is 30-40. The precipitation conditions therefore are relatively disadvantageous. This is further intensified by low humidity values, with an annual average of many years at 71-74%. Based on this data we may declare that the balance of precipitation and evaporation is negative in the region. The wetlands that have developed and exist can thank their subsistence to supplementary water influences (e.g. ground water).

The region so characteristic of the plain territories of the Carpathian Basin in terms of geology and climate offered settling opportunities for the natural continental fauna, including elements of both steppe and mediterranean nature. In addition, due to its geographical location (e.g. as a

basin centre surrounded by mountains) and its other features it became a place for the development of numerous endemic plant and animal species.

Under the given geological situation and under homogenous climate conditions the varied fauna is primarily the result of the abundance of pedological, hydrological and micro-relief conditions.

---

### General ecological features:

Lepidio-Puccinellietum and Astero-Agrostetum albae sodic marshes: The feature of this habitat is that it has only temporary or ephemeral saline water-flooding zone (0-10 cm), usually from later autumn to later spring. After it has dried out saline crystals often occur in high density on the bare surface, therefore, plant growth is strongly hampered here. The characteristic vegetation, which can thrive in these extreme conditions consists mainly of terrestrial halophyte and succulent plants such as: *Lepidion crassifolium*, *Puccinellia limosa*, *Camphorosma annua*, which occur sporadically on the surface. This habitat is one of the most important shorebird feeding zone because it has seasonal shallow (0-10 cm) water coverage and bare surface, often only the surface is wet and the muddy ground is soft. This feeding habitat exists mainly in spring and autumn for the waterbirds, when there is higher water level on the site.

Puccinellietum limosae sodic marshes: This habitat is similar as Lepidio marshes, but it has longer and a bit deeper saline water-flooding (0-20 cm), usually from early autumn to beginning of summer, due to more vegetation biomass can be found here, especially high dominant of halophyte *Puccinellia limosa*. The sodium crystal accumulation is not so expressed on the surface only in the deeper level of the soil. The vegetation coverage abundance may be very variable.

Bolboschoenus-Phragmitetum sodic marshes: This habitat regularly is covered with shallow water (0-30 cm) or wet all over the year. Due to here can overgrow more abundant halophyte vegetation as on saline marshes 2. zone. The characteristic dominant plants are *Bolboschoenus maritimus* and saline ecotype of *Phragmites communis*, these may occurrence in very different coverage proportion.

Open bed of pans: This habitat regularly is covered with deeper saline shallow water (10-50 cm) all over the year. According to there is not important emergent vegetation only few sporadic saline submerged water plants as *Potamogeton pectinatus*. This habitat is the major importance for waterbirds. In hot summer, when the water level may be dropped seriously, so pool-bed surfaces can be become only wet or dried out.

Other not characteristic wetlands types can be also found such as *Alopecuretum pratensis* meadow and *Caricetum acutiformis ripariae* marshes.

Continental Pannonic sodic affected steppes are extensively scattered around the wetlands such as *Artemisio-Festucetum pseudovinae danubiale*, and *Achilleo-Festucetum pseudovinae*. Fragmented Pannonic loess steppic grasslands are also such as *Salvio-Festucetum rupicola*.

---

**Bibliographical references:**

Andrikovics, S. & Murányi, D. (2003): Zoobentosz együttesekről a Szabadszállás-Fülpszállás környéki fehér szikes vizekben. With English summary: Zoobenthic assemblages in the white sodic waters near Szabadszállás-Fülöpszállás (Hungary). Természetvédelmi Közlemények 10. pp.251-271, 2003

Boros, E. (1996): Koegzisztens fészkelő partimadarak monitoringjának eredményei 1992-1995 között. With English summary: Preliminary results of the monitoring of coexistently breeding waders in Hungary between 1992 and 1995. Partimadár 5: 1-9p.

Boros, E. (1997): A Böddi-szék - egy kevésbé ismert alföldi szikes tó jellemzése a Bonaparte partfutó (*Calidris fuscicollis*) első előfordulása kapcsán. With English summary: First record of White-rumped Sandpiper in Hungary. Tűzok (Madártani Tájékoztató) 2.évf. 3.szám. 102-105.p.

Boros, E. (1998): Partimadár költőhabitatok térbeli makromintázatának vizsgálata két szikes tavon. With English summary: Spatial pattern of breeding habitats of three wader species at two saline lakes. Ornis Hungarica & Suppl. 1: 59-68. 1998

Boros, E. (1998): Kűszvágó csér (*Sterna hirundo*) fészkelőhelyek kialakítása a Kiskunsági Nemzeti Parkban. Partimadár. 6-7. (1997-1998): 1-8.

Boros, E. (1998): First record of White-rumped Sandpiper in Hungary., Aquila: vol. 103-104: p146-148. (1996-1997)

Boros, E. (1999): A magyarországi szikes tavak és vizek ökológiai értékelése. With English summary: Ecological state of sodic water bodies in Hungary. Acta Biol. Debr. Oecol. Hung. 9: p13-80 1999.

Boros, E.<sup>1</sup> & Biró, CS.<sup>2</sup> (1999): A Duna-Tisza közti szikes tavak ökológiai állapotváltozásai a XVIII-XX. századok időszakában. With English summary: Ecological change of sodic water bodies in the plain between Danube and Tisza from 18<sup>th</sup> to 20<sup>th</sup> centuries. Acta Biol. Debr. Oecol. Hung. 9: 81-105, 1999.

Boros, E. – Pigniczki, CS. (2001): Feltöltődött szikes tavak rekonstrukciója és a szikes mocsári vegetáció kezelése a kiskunsági szikes tavaknál. With English summary: Habitat reconstruction of charged sodic lakes and the maintain of the sodic marshland vegetation at the Kiskunsagian sodic lakes. Tűzok 6(1), 2001. p8-14.

Boros, E. (2002): Partimadarak (Charadrii) potenciális táplálékkínálata és az élőhelykezelés összefüggései kiskunsági szikes gyepterületeken. With English summary: Investigations on the relation between potential food resource of shorebird (Charadrii) and habitat management practices on sodic grassland habitats in the Kiskunság. Aquila, 107-108, p. 15-35.

Boros, E. (2003): Vízimadár populációváltozások és környezeti okai a Kiskunsági Nemzeti Park szikes tavain és mocsarain. With English summary: Waterbird population changes and their environmental reasons on sodic lakes of the Kiskunság National Park. Természetvédelmi Közlemények 10. pp. 289-312, 2003

Faragó, S. (1995): Geese in Hungary 1986-1991. IWRB Publication 36, 1995.

Faragó, S. (1996): A magyar vadlúd adatbázis 1984-1995: egy tartamos monitoring – Data base of gees in Hungary 1984-1995: A long-term monitoring. Magyar Vízivad Közlemények – Hungarian Waterfowl Publications No. 2. p3-222.

Fehér, G. (2003): Szikes területek vizeinek Desimales flórája. With English summary: Desimales-flora of sodic waters. Természetvédelmi Közlemények 10. pp.177-183, 2003

Fehér, G. & Schmidt, A. (2003): Kiskunsági szikes tavak (KNP II) algológiai vizsgálata. With English summary: Algological investigations of sodic ponds in Kiskunság. Természetvédelmi Közlemények 10. pp.163-176, 2003

Forró, L. (2003): A Fülöpszállás-Szabadszállás környéki szikes vizek rákfaunája. With English summary: The microcrustacean fauna of the sodic waters in the vicinity of Fülöpszállás-Szabadszállás. Természetvédelmi Közlemények 10. pp.241-250, 2003

Forró, L. & Boros, E. (1997): Microcrustacean zooplankton as potential food of *Recurvirostra avosetta* in sodic waters of the Hungarian Plain. Lymnology and Waterfowl monitoring, modelling and management. Hungarian Waterfowl Publications No.3. (Ed. S. Faragó and J. Kerekes) Sarród-Sopron, 21-23. nov. 1994. 239-250p.

Füzi, A. – Biró, B. & Tóth, T. (2003): Növény-mikróba kölcsönhatások és néhány talajtulajdonság közötti összefüggés hazai szikeseken. With English summary: Plant-microba interactions and the effect of some soil properties on salt affected areas. Természetvédelmi Közlemények 10. pp.191-205, 2003

Mahunka, S. [eds.] 1986: The fauna of the Kiskunság National Park, Vol I. Akadémia Kiadó, Budapest 1986.

Mahunka, S. [eds.] 1987: The fauna of the Kiskunság National Park, Vol II. Akadémia Kiadó, Budapest 1987.

Papp, B. (2003): A Fülöpszállás környéki szikes tavak mohafiórájának állapotfelmérése. With English summary: Investigations on the bryophyte flora of saline-alkali lakes around Fülöpszállás. Természetvédelmi Közlemények 10. pp.217-228, 2003

Pálfai, I. (1994), Összefoglaló tanulmány a Duna–Tisza közti talajvízszint-süllyedés okairól, és a vízhiányos helyzet javításának lehetőségeiről. – In: Pálfai I. (szerk.) 1994: A Nagyalföld Alapítvány Kötetei 3. A Duna–Tisza közti hátság vízgazdálkodási problémái. – Nagyalföld Alapítvány, Békéscsaba. pp. 111-126. ISBN 963 04 3942 1, ISSN 1216-4933.

Schmidt, A (2003): Kiskunsági szikes tavak (KNP II) összehasonlító vízkémiai vizsgálata. With English summary: Comparing water chemical investigations of sodic ponds in the Kiskunság National Park (KNP II). Természetvédelmi Közlemények 10. pp.153-162, 2003

Szabó, A. (2003): Protozoológiai (Protozoa, Ciliata) vizsgálatok a Fülöpszállás-Szabadszállás környéki szikes kisvizekben. With English summary: Recent protozoological investigations of small alkaline wetland patches in Fülöpszállás-Szabadszállás area. Természetvédelmi Közlemények 10. pp.229-239, 2003

Szabó, S. 1993: The effect of becoming waterless and experiments of livingplace reconstruction on Mollusca living in the soid laces of Upper Kiskunság. *Malacological Newsletters*. 12. 47-57p.

Szabó, S. 1990: The survival, resettle and Return of the protected snails living in the National Parks of Hungary, - Abstr. 2<sup>nd</sup> Int. Congr. on Medical and Applied Malacology (Seul, Korea 1990

Szabó, S. (2003): Két évtized a Felső-Kiskunság szikes vizeiben élő Molluscák kutatásában. With English summary: Aquatic Mollusca of the sodic waters in the Upper Kiskunság. *Természetvédelmi Közlemények* 10. pp.273-288, 2003

Szujkó-Lacza, J. & Kováts, D. (eds.) 1993: The Flora of the Kiskunság National PArk. In the Danube-Tisza Mid-Region of Hungary. Vol. I. Magyar Természettudományi Múzeum, Budapest 1993. 469pp.

Sterbetz, I. 1981. Protected wetlands of international importance in Hungary. IWRB XVII. International Conference in Debrecen 1981.

Tóth, T. – Kuti, L. & Fügedi, U. (2003): Havonkénti vizsgálatok a Zab-szék mellett. A tóvíz, a talajvíz, a talaj, növényzet időbeli változásai. With English summary: Monthly observations at Zab-szék saline lake. Temporal changes of lake water and groundwater, soils and vegetation. *Természetvédelmi Közlemények* 10. pp.191-205, 2003

Vörös, L. & Balogh, K. (2003): Fotoautrotóf pikoplankton Duna-Tisza közti szikes tavakban. With English summary: Photoautotrophic picoplankton of the natron lakes in the Danube-Tisza Interfluve. *Természetvédelmi Közlemények* 10. pp.185-189, 2003