Information Sheet on Ramsar Wetlands (RIS)

Categories approved by Recommendation 4.7, as amended by Resolution VIII.13 of the Conference of the Contracting Parties.

Note for compilers:

- The RIS should be completed in accordance with the attached Explanatory Notes and Guidelines for completing the Information Sheet on Ramsar Wetlands. Compilers are strongly advised to read this guidance before filling in the RIS.
- Once completed, the RIS (and accompanying map(s)) should be submitted to the Ramsar Secretariat. Compilers are

strongly urged to provide an electronic (MS Word) copy of the RIS a	and, where possible, digital copies of maps.
1. Name and address of the compiler of this form:	FOR OFFICE USE ONLY.
	DD MM YY
Joint Nature Conservation Committee	
Monkstone House	
City Road	Designation date Site Reference Number
Peterborough	
Cambridgeshire PE1 1JY	
UK	
Telephone/Fax: +44 (0)1733 - 562 626 / +44 (0)17	733 – 555 948
Email: <u>RIS@JNCC.gov.uk</u>	
2 Date this sheet was completed/undeted	
2. Date this sheet was completed/updated:	
09 March 2006	
3. Country:	
UK (Scotland)	
4. Name of the Ramsar site:	
South Uist Machair and Lochs	
	_
5. Map of site included:	
Refer to Annex III of the Explanatory Notes and Guidelines, for detailed g	guidance on provision of suitable maps.
a) hard copy (required for inclusion of site in the Ramsar Li	ist): yes ✓ -or- no
b) digital (electronic) format (optional): Yes	
6. Geographical coordinates (latitude/longitude):	
57 18 50 N 07 20 00 W	
7. General location:	
Include in which part of the country and which large administrative region	n(s), and the location of the nearest large town.
Nearest town/city: Benbecula	
South Uist Machair and Lochs, which includes Lochs Druidi	ibeg, a' Machair and Stilligarry, lies on the
west coast of South Uist in the Outer Hebrides.	
Administrative region: Western Isles / Na h-Eileanan an Iar	r
8. Elevation (average and/or max. & min.) (metres): 9.	Area (hectares): 5019.05
Min1	Alea (nectates). 3019.03
Max. 38	
Mean 6	
1110411	

Ramsar Information Sheet: UK13058 Page 1 of 9	South Uist Machair and Lochs
---	------------------------------

10. Overview:

Provide a short paragraph giving a summary description of the principal ecological characteristics and importance of the

South Uist Machair and Lochs Ramsar site is a complex site along the west coast of South Uist. This area is of outstanding importance for its transition of habitats from acidic moorland to the calcareous coastal plain, and for the transition from freshwater habitats to saltwater habitats. This complex includes outstanding examples of, moving seawards, relict woodland, moorland and blanket bog, large oligotrophic lochs, acidic blacklands, wet and dry machair with eutrophic machair lochs, freshwater marsh, saltmarsh, coastal dunes and sandy rocky shores. These areas are of outstanding importance for their populations of wintering and breeding waterfowl.

11. Ramsar Criteria:

Circle or underline each Criterion applied to the designation of the Ramsar site. See Annex II of the Explanatory Notes and Guidelines for the Criteria and guidelines for their application (adopted by Resolution VII.11).

1, 2, 6

Secretariat comment: The RIS provides information requiring the application of Criterion 4. This needs to be included in the next update.

12. Justification for the application of each Criterion listed in 11. above:

Provide justification for each Criterion in turn, clearly identifying to which Criterion the justification applies (see Annex II for guidance on acceptable forms of justification).

Ramsar criterion 1

The site contains some of the finest examples of a continual gradient from marine to freshwater habitats and is of outstanding biological and geomorphological significance through containing the largest machair system in Britain.

Ramsar criterion 2

The site supports an assemblage of rare wetland plants and animals. The flora includes at least four Nationally Scarce vascular plants, slender naiad Najas flexilis, cowbane Cicuta virosa, slender-leaved pondweed Potamogeton filiformis and Baltic rush Juncus balticus. The fauna includes otter Lutra lutra and salmon Salmo salar.

Ramsar criterion 6 – species/populations occurring at levels of international importance.

Qualifying Species/populations (as identified at designation):

Species regularly supported during the breeding season:

Dunlin, Calidris alpina schinzii, 382 pairs, representing an average of 3.5% of the Baltic/UK/Ireland breeding population (2000)

Species with peak counts in winter:

Ringed plover, Charadrius hiaticula, 66 individuals, representing an average of 0.2% Europe/Northwest Africa of the GB population (5 year peak mean 1998/9-

2002/3)

Contemporary data and information on waterbird trends at this site and their regional (sub-national) and national contexts can be found in the Wetland Bird Survey report, which is updated annually. See www.bto.org/survey/webs/webs-alerts-index.htm.

13. Biogeography (required when Criteria 1 and/or 3 and /or certain applications of Criterion 2 are applied to the designation):

Name the relevant biogeographic region that includes the Ramsar site, and identify the biogeographic regionalisation system that has been applied.

a) biogeographic region:

Atlantic

b) biogeographic regionalisation scheme (include reference citation):

Council Directive 92/43/EEC

14. Physical features of the site:

Describe, as appropriate, the geology, geomorphology; origins - natural or artificial; hydrology; soil type; water quality; water depth, water permanence; fluctuations in water level; tidal variations; downstream area; general climate, etc.

Soil & geology	acidic, basic, shingle, sand, peat, nutrient-rich, nutrient-	
	poor, metamorphic, gravel, pebble	
Geomorphology and landscape	lowland, island, coastal, slope, crags/ledges, subtidal	
	sediments (including sandbank/mudbank), intertidal	
	sediments (including sandflat/mudflat), open coast	
	(including bay), pools, intertidal rock	
Nutrient status	eutrophic, mesotrophic, oligotrophic	
pН	acidic, alkaline, circumneutral	
Salinity	brackish / mixosaline, fresh, saline / euhaline	
Soil	mainly mineral	
Water permanence	usually permanent	
Summary of main climatic features	Annual averages (Stornoway, 1971–2000)	
	(www.metoffice.com/climate/uk/averages/19712000/sites	
	/stornoway.html)	
	Max. daily temperature: 11.0° C	
	Min. daily temperature: 5.6° C	
	Days of air frost: 30.7	
	Rainfall: 1231.7 mm	
	Hrs. of sunshine: 1216.8	

General description of the Physical Features:

South Uist Machair and Lochs is a complex site of outstanding importance for its transition of habitats from acidic moorland to the calcareous coastal plain, and for the transition from freshwater to saltwater habitats. This complex includes outstanding examples of relict woodland, moorland and blanket bog, large oligotrophic lochs, acidic blacklands, wet and dry machair with eutrophic machair lochs, freshwater marsh, saltmarsh, coastal dunes and sandy and rocky shores.

15. Physical features of the catchment area:

Describe the surface area, general geology and geomorphological features, general soil types, general land use, and climate (including climate type).

South Uist Machair and Lochs is a complex site of outstanding importance for its transition of habitats from acidic moorland to the calcareous coastal plain, and for the transition from freshwater to saltwater habitats. This complex includes outstanding examples of relict woodland, moorland and blanket bog, large oligotrophic lochs, acidic blacklands, wet and dry machair with eutrophic machair lochs, freshwater marsh, saltmarsh, coastal dunes and sandy and rocky shores.

16. Hydrological values:

Describe the functions and values of the wetland in groundwater recharge, flood control, sediment trapping, shoreline stabilization, etc.

No special values known

17. Wetland types

Inland wetland, Marine/coastal wetland

Code	Name	% Area
Е	Sand / shingle shores (including dune systems)	34.8
F	Estuarine waters	0.2
О	Freshwater lakes: permanent	12
Q	Saline / brackish lakes: permanent	14.9
Тр	Freshwater marshes / pools: permanent	11.9
U	Peatlands (including peat bogs swamps, fens)	13.4
Other	Other	12.8

18. General ecological features:

Provide further description, as appropriate, of the main habitats, vegetation types, plant and animal communities present in the Ramsar site.

South Uist Machair & Lochs is a complex site along the west coast of South Uist, which is of outstanding importance for its transition of wetland habitats from blanket bog to calcareous coastal plain, and for the transition through freshwater to saltwater habitats. The site has outstanding examples of blanket bog, large oligotrophic lochs, acidic blacklands, wet and dry machair with machair lochs, freshwater marsh, saltmarsh, coastal dunes and sandy and rocky shores.

The machair areas include wet and dry machair, some of which is uncultivated. Together with the associated marsh, fen and eutrophic machair lochs the coastal plain supports a rich and diverse range of plant communities, including some locally or nationally scarce plants. The estuarine and brackish water habitats of the Howmore River, Loch Roag and the freshwater Loch Fada show one of the best examples of gradation from estuarine to freshwater habitats, along with associated gradation of flora and fauna, in Great Britain. Loch Bee is the largest brackish loch in the Western Isles and exhibits a gradation in salinity from freshwater to seawater. The coast, machair, lochs and blackland are of considerable ornithological importance for wintering and breeding birds.

19. Noteworthy flora:

Provide additional information on particular species and why they are noteworthy (expanding as necessary on information provided in **12**. Justification for the application of the Criteria) indicating, e.g. which species/communities are unique, rare, endangered or biogeographically important, etc. *Do not include here taxonomic lists of species present – these may be supplied as supplementary information to the RIS.*

Assemblage.

The site is internationally important because it supports the following Habitats Directive Annex I features:

H1150	Coastal lagoons
H1210	Annual vegetation of drift lines
H2120	Shifting dunes along the shoreline with <i>Ammophila arenaria</i> ("white dunes")
H2130	Fixed dunes with herbaceous vegetation ("grey dunes")
H2190	Humid dune slacks
H21A0	Machairs
H3110	Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia
uniflorae)	
H3130	Oligotrophic to mesotrophic standing waters with vegetation of the <i>Littorelletea</i>

H3130 Oligotrophic to mesotrophic standing waters with vegetation of the *Littorelletea* uniflorae and/or of the *Isoëto-Nanojuncetea*

H3140 Hard oligo-mesotrophic waters with benthic vegetation of *Chara* spp.

H3150 Natural eutrophic lakes with *Magnopotamion* or *Hydrocharition*-type vegetation

Internationally important species occurring on the site:

S1833 Najas flexilis Slender naiad (Habitats Directive Annex II species)

Nationally important species occurring on the site:

Higher Plants.

Potamogeton filiformis, Cicuta virosa, Juncus baltica

Lower Plants.

Chara spp.

20. Noteworthy fauna:

Provide additional information on particular species and why they are noteworthy (expanding as necessary on information provided in **12**. Justification for the application of the Criteria) indicating, e.g. which species/communities are unique, rare, endangered or biogeographically important, etc., including count data. *Do not include here taxonomic lists of species present* – these may be supplied as supplementary information to the RIS.

Rirde

Species currently occurring at levels of national importance:

Species regularly supported during the breeding season:

Corn crake, *Crex crex*, Europe 15 individuals, representing an average of 2.5% of

the GB population (5 year mean 1993-1997)

Little tern, Sterna albifrons albifrons, W Europe 24 apparently occupied nests, representing an

average of 1.2% of the GB population (Seabird

393 pairs, representing an average of 2.5% of the

2000 Census)

Ringed plover, *Charadrius hiaticula hiaticula*, breeding in southern Scandinavia/Baltic, British Isles, N & E continental Europe; wintering in

biogeographic breeding population (Count as at 1995)

Europe, Med. & N Africa

Species Information

Internationally important species occurring on the site.

S1355 Lutra lutra Otter (Habitats Directive Annex II species)

Nationally important species occurring on the site.

Invertebrates.

Bombus distinguendus, Lycia zonaria, Colletes floralis, Thanatophilus dispar

21. Social and cultural values:

e.g. fisheries production, forestry, religious importance, archaeological sites, social relations with the wetland, etc. Distinguish between historical/archaeological/religious significance and current socio-economic values.

Aesthetic

Aquatic vegetation (e.g. reeds, willows, seaweed)

Archaeological/historical site

Environmental education/interpretation

Livestock grazing

Non-consumptive recreation

Scientific research

Sport fishing

Sport hunting

Tourism

Traditional cultural

22. Land tenure/ownership:

Ownership category	On-site	Off-site
National/Crown Estate	+	
Private	+	+
Public/communal	+	

23. Current land (including water) use:

Activity	On-site	Off-site
Nature conservation	+	+
Tourism	+	+
Recreation	+	+
Collection of non-timber natural	+	+
products: commercial		
Collection of non-timber natural	+	+
products: subsistence		
Cutting of vegetation (small-	+	+
scale/subsistence)		
Fishing: commercial	+	+
Fishing: recreational/sport	+	
Shifting arable agriculture	+	
Rough or shifting grazing	+	+
Permanent pastoral agriculture	+	+
Hay meadows	+	+
Hunting: recreational/sport	+	
Sewage treatment/disposal	+	+
Mineral exploration (excl.	+	
hydrocarbons)		
Non-urbanised settlements	+	+
Military activities	+	

24. Factors adversely affecting the site's ecological character, including changes in land (including water) use and development projects:

Explanation of reporting category:

- 1. Those factors that are still operating, but it is unclear if they are under control, as there is a lag in showing the management or regulatory regime to be successful.
- 2. Those factors that are not currently being managed, or where the regulatory regime appears to have been ineffective so far.

NA = Not Applicable because no factors have been reported.

Adverse Factor Category	Reporting Category	Description of the problem (Newly reported Factors only)	On-Site	Off-Site	Major Impact?
Introduction/invasion of non-native animal species	1		+	+	+

For category 2 factors only.
What measures have been taken / are planned / regulatory processes invoked, to mitigate the effect of these factors'

Is the site subject to adverse ecological change? YES

25. Conservation measures taken:

List national category and legal status of protected areas, including boundary relationships with the Ramsar site; management practices; whether an officially approved management plan exists and whether it is being implemented.

Conservation measure	On-site	Off-site
Site/ Area of Special Scientific Interest	+	
(SSSI/ASSI)		
Special Protection Area (SPA)	+	
Site management statement/plan implemented	+	
Other	+	
Special Area of Conservation (SAC)	+	

26. Conservation measures proposed but not yet implemented:

e.g. management plan in preparation; official proposal as a legally protected area, etc.

No information available

27. Current scientific research and facilities:

e.g. details of current research projects, including biodiversity monitoring; existence of a field research station, etc.

Fauna.

Research on effects of hedgehogs on breeding waders. Corncrake surveys (RSPB) annual.

Greylag goose breeding monitoring on Loch Druidibeg NNR.

Invertebrates: Survey of *Bombus distinguendus* in 1997 and ongoing.

Birds: Breeding wader surveys in 1983, 1993 and 2000, wintering wader survey 1994-95 and 2004.

Various national British Trust for Ornithology surveys.

Flora.

Trial rabbit exclusion zones to gauge effects of grazing and burrowing on machair.

Freshwater vegetation surveyed in 1995. Machair vegetation surveyed 1995, 1988. Machair site condition monitoring (SCM) 2004. Lagoon SCM 2002.

28. Current conservation education:

e.g. visitor centre, observation hides and nature trails, information booklets, facilities for school visits, etc.

There are interpretive panels, a reserve leaflet, and a limited number of guided walks for visitors and organised parties on the Loch Druidibeg NNR section of the site.

29. Current recreation and tourism:

State if the wetland is used for recreation/tourism; indicate type(s) and their frequency/intensity.

Activities, Facilities provided and Seasonality.

Land-based recreation:

Walking, including dog-walking, birdwatching and mammal watching occur, at a fairly low level, over much of the site, all year. The south-western corner of Loch Druidibeg has restricted access during the spring and summer, to reduce disturbance to breeding birds.

South Uist Estates shoot wildfowl over the part of the site they own. The season is from 1 September to 12 February inclusive.

Water-based recreation:

Shooting season 12 August-20 February.

Angling occurs from both shore and boat. Again this mainly occurs from April to September. It is restricted on Loch Druidibeg and no boats are allowed there.

30. Jurisdiction:

Include territorial, e.g. state/region, and functional/sectoral, e.g. Dept. of Agriculture/Dept. of Environment, etc.

Scottish Executive, Environment and Rural Affairs Department

31. Management authority:

Provide the name and address of the local office(s) of the agency(ies) or organisation(s) directly responsible for managing the wetland. Wherever possible provide also the title and/or name of the person or persons in this office with responsibility for the wetland.

Scottish Natural Heritage, 2 Anderson Place, Edinburgh, EH6 5NP

32. Bibliographical references:

Scientific/technical references only. If biogeographic regionalisation scheme applied (see 13 above), list full reference citation for the scheme.

Site-relevant references

- Barne, JH, Robson, CF, Kaznowska, SS, Doody, JP, Davidson, NC & Buck, AL (eds.) (1997) *Coasts and seas of the United Kingdom. Regions 15 & 16. North-west Scotland: the Western Isles and west Highland.* Joint Nature Conservation Committee, Peterborough. (Coastal Directories Series.)
- Booth, AB & Milne, F (1999) Western Isles corncrake habitat survey. (Contactor: Central Environmental Surveys.) Scottish Natural Heritage Commissioned Report, No. F99LC03
- Boyd, JM (ed.) (1979) The natural environment of the Outer Hebrides. Proceedings of the Royal Society of Edinburgh. Series B: Biological Sciences, 77
- Boyd, JM & Boyd, IL (1990) The Hebrides. A natural history. Collins, London (New Naturalist No. 76)
- Bryan, A (1994) The Minch review. Scottish Natural Heritage, Stornoway
- Connor, DW & Little, M (1998) Chapter 14. Outer Hebrides (MNCR Sector 14). In: *Benthic marine ecosystems of Great Britain and the north-east Atlantic*, ed. by K. Hiscock, 371-383. Joint Nature Conservation Committee, Peterborough. (Coasts and Seas of the United Kingdom. MNCR series)
- Cranswick, PA, Waters, RJ, Musgrove, AJ & Pollitt, MS (1997) *The Wetland Bird Survey 1995–96: wildfowl and wader counts.* British Trust for Ornithology, Wildfowl and Wetlands Trust, Royal Society for the Protection of Birds & Joint Nature Conservation Committee, Slimbridge
- Dargie, TCD (1993) Sand dune vegetation survey of Great Britain: a national inventory. Part II: Scotland. Joint Nature Conservation Committee, Peterborough
- Dargie, TCD (2000) Sand dune vegetation survey of Scotland: national report. *Scottish Natural Heritage, Commissioned Report*, No. F97AA401. www.snh.org.uk/pdfs/strategy/commreports/F97AA401.pdf
- Ecology UK Ltd. (2005) North Uist Machair and Islands SPA and South Uist Machair and Lochs SPA: winter wader survey 2004. Scottish Natural Heritage Commissioned Report, No. **097** (ROAME No. F03LC05)
- Johnston, R & Mudge, G (1996) South Uist Machair & Lochs Proposed SPA/RAMSAR Site (108A), Departmental brief. Scottish Natural Heritage, Edinburgh (International Sites Series)
- Johnston, R, Boyle, J & Quick, R (1995) A survey of wintering waders on the west coast of the Uists in winter 1994–95. Scottish Natural Heritage, Edinburgh (unpublished report)
- Johnston, R, Murray, J & Shepherd, MA (1995) Survey of breeding waders on Uist SSSIs. Scottish Natural Heritage, Edinburgh
- May, VJ & Hansom, JD (eds.) (2003) Coastal geomorphology of Great Britain. Joint Nature Conservation Committee, Peterborough (Geological Conservation Review Series, No. 28)
- McLeod, CR, Yeo, M, Brown, AE, Burn, AJ, Hopkins, JJ & Way, SF (eds.) (2004) *The Habitats Directive: selection of Special Areas of Conservation in the UK*. 2nd edn. Joint Nature Conservation Committee, Peterborough. www.jncc.gov.uk/SACselection
- Ramsay, DL & Brampton, AH (2000) Coastal cells in Scotland: Cells 8 & 9 The Western Isles. Scottish Natural Heritage Research Survey and Monitoring Report, No. 150
- Ratcliffe, DA (ed.) (1977) A Nature Conservation Review. The selection of biological sites of national importance to nature conservation in Britain. Cambridge University Press (for the Natural Environment Research Council and the Nature Conservancy Council), Cambridge (2 vols.)
- Ritchie, W (1966) The physiography of the machair of South Uist. Unpublished PhD thesis, University of Glasgow
- Ritchie, W (1969) *The coastal geomorphology of North Uist*. University of Aberdeen, Department of Geography (O'Dell Memorial Monographs, No. 1)
- Ritchie, W (1971) The beaches of Barra and the Uists. A survey of the beach, dune and machair areas of Barra, South Uist, North Uist and Berneray. University of Aberdeen, Department of Geography, for Countryside Commission for Scotland, Perth. [Reprinted 2004 as Scottish Natural Heritage Commissioned Report, No. **047**]
- Scottish Natural Heritage (199*) Loch Druidibeg NNR Management Plan 1995-2000. Scottish Natural Heritage South Uist Office, Stilligarry
- Scottish Natural Heritage (1998) Machair. Scottish Natural Heritage, Edinburgh (Scotland's Living Landscapes)
- Scottish Natural Heritage (2004) *Uist Wader Project webpages*. Scottish Natural Heritage, Edinburgh. www.snh.org.uk/index/i-frame.htm
- Sullivan, C (1995) The carrion beetle *Thanatophilus dispar* at Loch Druidibeg NNR. *Scottish Natural Heritage Commissioned Report*, No. **NW593** (unpublished report)

Information Sheet on Ramsar Wetlands (RIS), page 9

- Stewart, NF (2004) Important stonewort areas. An assessment of the best areas for stoneworts in the United Kingdom. Plantlife International, Salisbury
- Stroud, DA, Chambers, D, Cook, S, Buxton, N, Fraser, B, Clement, P, Lewis, P, McLean, I, Baker, H & Whitehead, S (eds.) (2001) *The UK SPA network: its scope and content*. Joint Nature Conservation Committee, Peterborough (3 vols.) www.jncc.gov.uk/UKSPA/default.htm
- Thorpe, K, Dalkin, MJ, Fortune, F & Nichols, DM (1998) Marine Nature Conservation Review Sector 14. Lagoons in the Outer Hebrides: area summaries. Joint Nature Conservation Committee, Peterborough (Coasts and seas of the United Kingdom. MNCR series)
- Weighell, AJ, Donnelly, AP & Calder, K (eds.) (2000) *Directory of the Celtic coasts and seas*. Joint Nature Conservation Committee, Peterborough

Please return to: Ramsar Secretariat, Rue Mauverney 28, CH-1196 Gland, Switzerland Telephone: +41 22 999 0170 • Fax: +41 22 999 0169 • email: ramsar@ramsar.org

Ramsar Information Sheet: UK13058 Page 9 of 9 South Uist Machair and Lochs