Color codes

Fields back-shaded in light blue relate to data and information required only for RIS updates.

Note that some fields concerning aspects of Part 3, the Ecological Character Description of the RIS (tinted in purple), are not expected to be completed as part of a standard RIS, but are included for completeness so as to provide the requested consistency between the RIS and the format of a 'full' Ecological Character Description, as adopted in Resolution X.15 (2008). If a Contracting Party does have information available that is relevant to these fields (for example from a national format Ecological Character Description) it may, if it wishes to, include information in these additional fields.

1 - Summary

Summary

Lake Nakuru is a shallow, strongly alkaline lake, with surrounding woodland and grassland set in a picturesque landscape within the Lake Nakuru National Park that abuts Nakuru town. The lake is located in a closed basin of about 1800 km2 of which 3,300 ha forms the main lake water body. The lake's high alkalinity, conductivity and other physical - chemical parameters makes it uninhabitable to many aquatic species; however the few that have adapted to these harsh conditions have made the lake to have one of the highest producer biomass among Kenya's southern Rift Valley alkaline lakes. The foundation of the lake's simple food chains is the cyanophyte Arthrospira fusiformis, which often occurs as unialgal bloom. At such times, it can support huge numbers of Lesser Flamingo (Phoeniconaias minor). The introduction of fish species Oreochromis alcalicus grahami in 1960s has attracted a number of other secondary consumers including several piscivorous bird species. The lake's catchments has experienced expansion in agriculture and urban developments. There has been accidental introduction of nile tilapia Oreochromis nilotica. Increased numbers of fresh water fish eating bird species has been recorded. In the recent past, the lake has expanded in volume and height due high precipitation attributed to climate change, this has resulted in a drastic decline in flamingo numbers recorded.

2 - Data & location

2.1 - Formal data

2.1.1 - Name and address of the com	piler of this RIS
Responsible compiler	
Institution/agency	Wildlife Research and Training Institute
Postal address	842-20117
National Ramsar Administrati	ve Authority
Institution/agency	Kenya Wildlife Service
Postal address	P.O. Box 40241 – 00100 Nairobi, Kenya
2.1.2 - Period of collection of data and	d information used to compile the RIS
From year	2016
To year	2022
2.1.3 - Name of the Ramsar Site	
Official name (in English, French or Spanish)	Lake Nakuru
Unofficial name (optional)	Lake Nakuru National Park
	d area of the Site since its designation or earlier update
	Changes to Site boundary Yes O No ⊚
(Update) For secretariat only: TI	B. Changes to Site area No change to area
2.1.5 - Changes to the ecological cha (Update) 6b i. Has the ecological character of tapplicable Criteria) change	racter of the Site the Ramsar Site (including Yes (likely)
аррисавле Сптепа) спанув	(Update) Are the changes Positive O Negative O Positive & Negative ●
(Update) Positive %	50
(Update) Negative %	50
(Update	a) No information available
(Update) Optional text box to provide further info	rmation
Excessive flooding since 2010 affect observed in the years.	ting habitat suitability for waterbirds especially the lesser flamingos with sporadic to decline in sightings
(Update) Changes resulting from causes op	perating within the existing boundaries?
(Update) Changes resulting from causes o	pperating beyond the site's boundaries?
(Update) Changes consequent upon site bounthe exclusion of some wetland types formerly	
(Update) Changes consequent upon site bour the inclusion of different	ndary increase alone (e.g., wetland types in the site)?
	cological character of the Ramsar Site, including in the application of the Criteria, since the previous RIS for the site.
Excessive flooding affecting habitat	suitability for waterbirds especialky flamingos. Sporadic tono sightings observed in the years. Critarion5.
(Update) Is the change in ecological character AND a significant change (above the li	negative, human-induced minit of acceptable change)

2.2 - Site location

2.2.1 - Defining the Site boundaries

b) Digital map/image

<3 file(s) uploaded?

Earmar mana			
	U		
. omioi mapo			

Boundaries description

The site is located in the Eastern Rift Valley in Kenya. It follows the designated and gazetted boundary of Lake Nakuru National Park which was gazetted as a wildlife protected area in 1974. It is surrounded by Nakuru City and settlement to the North, Eastern and Western Boundaries. The southern Boundary is connected to Soysambu Conservancy.

2.2.2 - General location

a) In which large administrative region does the site lie?	Nakuru County
tile site lie:	
b) What is the nearest town or population	Nakuru City

2.2.3 - For wetlands on national boundaries only

a) Does the wetland extend onto the territory of one or more other countries? Yes O No \odot

b) Is the site adjacent to another designated Ramsar Site on the territory of another Contracting Party?

2.2.4 - Area of the Site

Official area, in hectares (ha): 18800

Area, in hectares (ha) as calculated from 19035.758

2.2.5 - Biogeography

Biogeographic regions

Diogeographic regions	
Regionalisation scheme(s)	Biogeographic region
WWF Terrestrial Ecoregions	East African Halophytics

Other biogeographic regionalisation scheme

Specify the biogeographic regionalisation scheme and the biogeographic region please

3 - Why is the Site important?

3.1 - Ramsar Criteria and their justification

<no data available>

Criterion 2 : Rare species and threatened ecological communities

Optional text box to provide further

Lake Nakuru is globally renowned for hosting the largest population of Lesser Flamingos (Near threatened)—what has been described as "the world's greatest ornithological spectacle". With over 450 species of birds, 50 species of mammals and 300 species of plants, it is one of the highest biodiversity parks in the country. The lake and its littoral area support over 70 species of waterfowl and water-related birds. Each year the resident bird life of the park is enriched by the presence of several species of Palearctic waders that use the lake as a staging ground during their winter migration down the Rift Valley fly way (GoK and UNDP, 2021)

Criterion 3 : Biological diversity

Lake Nakuru supports over 450 avian species and about 70 waterbird species, including 1 million flamingos. The lake also supports large densities of cynophytes Spirulina platensis, characteristic of the Rift Valley saline lakes. The spirulina in turn supports sodic fish Tilapia grahami (Oreochromis alcalicus grahami) introduced into the lake from Lake Magadi in the 1960s. Introduction of this fish has enriched waterbird diversity by supporting hundreds of fish eating water birds like Pelicans and Cormorants.

Criterion 4 : Support during critical life cycle stage or in adverse conditions

Optional text box to provide further

Lake Nakuru offers display and refuge site to the near threatened Lesser Flamingo (Phoeniconaias minor) and Greater Flamingo (Phoenicopterus ruber roseus). During water bird counts conducted in January 2021 a total of 19602 individual birds were counted falling in 77 species (Irene et.al 2021). Breeding attempts for Lesser Flamingo has also been reported from the Lake. The Lake is a pathway and staging site for several Palearctic migrants during migration seasons. The springs and their associated wetlands found at the Lake is a source of drinking water for birds and mammals during critical breeding stages.

☑ Criterion 5 : >20,000 waterbirds

Overall waterbird numbers 25.551

Start year 2021

2021 End year

Source of data: The National Water Bird Census for Kenya Report, 2021 & 2023

Optional text box to provide further

Over 77 species of birds have been identified at Lake Nakuru including during the early and mid - year water bird counts. The Lake supports over 20,000 water birds for different species of water birds. In the year 2021, 25,551 individual birds were counted while 19,299 individuals were counted in 2023. The number of birds has been fluctuating due to changes in water chemistry of the lake.

☑ Criterion 6 : >1% waterbird population

Optional text box to provide further information

Previously, in the five - year period of bird counts (1999 - 2003) Lake Nakurusupported over 1% of the East African Flamingo populations of Lesser Flamingo (Phoeniconaias minor) (average of 488,908 birds; 24.4%) and Greater Flamingos (Phoenicopterus ruber roseus) (average of 3,519 birds; 10.05%). Now with the increasing water levels, the large numbers of flamingos have since disappeared to other rift valley lakes

3.2 - Plant species whose presence relates to the international importance of the site

<no data available>

3.3 - Animal species whose presence relates to the international importance of the site

Phylum	Scientific name	Species qualifies under criterion	co	Species ontributes under criterion		Period of pop. Est.	0/_	III CNI		CMS Appendix I	Other Status	Justification
Others												
CHORDATA / ACTINOPTERYGII	Alcolapia grahami		\checkmark					VU				
CHORDATA / MAMMALIA	Ceratotherium simum		V					NT	1			
CHORDATA / MAMMALIA	Diceros bicornis							CR	\mathscr{L}			
CHORDATA/ MAMMALIA	Giraffa camelopardalis		V					VU			Endangered	population decreasing
CHORDATA/ MAMMALIA	Hippopotamus amphibius							VU				
CHORDATA / MAMMALIA	Panthera leo							VU	\mathscr{L}			
CHORDATA/ MAMMALIA	Panthera pardus		V					VU	\checkmark		current status Near threatened	
Birds										<u>'</u>		
CHORDATA / AVES	Anhinga rufa		V					LC			regionally threatened	
CHORDATA / AVES	Aquila clanga		V					VU		V	Endangered	A migrant species
CHORDATA / AVES	Aquila heliaca		V					VU	✓	V		Amigrant
CHORDATA / AVES	Ardea alba		V					LC			regionally threatened	
CHORDATA / AVES	Ardeola idae							EN		4		
CHORDATA / AVES	Balearica regulorum							EN				
CHORDATA / AVES	Falco naumanni							LC		V		
CHORDATA / AVES	Phoeniconaias minor				10437	2021	0.52	NT			CITES Appendix II	display and refuge site
CHORDATA / AVES	Phoenicopterus roseus		V		446	2021	0.46	LC			CITES Appendix II	display and refuge site
CHORDATA / AVES	Podiceps cristatus		V					LC			regionally threatened	
CHORDATA / AVES	Prionops poliolophus		V					NT			regionally threatened	

¹⁾ Percentage of the total biogeographic population at the site

Lake Nakuru offers display and refuge site to the threatened Lesser Flamingo (Phoeniconaias minor) and Greater Flamingo (Phoenicopterus ruber roseus). During water bird counts conducted at the Lake since 1990s, a significant percentage of juvenile flamingos have been counted at the lake. Breeding attempts for Lesser Flamingo has also been reported from the Lake. The Lake is a pathway and staging site for several Palearctic migrants during migration seasons. The springs and their associated wetlands found at the Lake is a source of drinking water for birds and mammals during critical breeding stages.

3.4 - Ecological communities whose presence relates to the international importance of the site

Name of ecological community	Community qualifies under Criterion 2?	Description	Justification
Planktons	Ø	Diatoms and other algae that bloom occasionally form substantial amount of total biomass	Food for the flamingo and fish
Algae	Ø	The main primary producer of the open water zone is the blue - green algae Spirulina platensis that constitutes 95% of the algal population	The blue green algae is the primary food for the lesser flamingo
Diatoms	2	Diatoms and other algae that bloom occasionally form substantial amount of total biomass	Contitute part of the lake's total biomass

Optional text box to provide further information

Diatoms and other algae that bloom occasionally form substantial amount of total biomass. The main primary producer of the open water zone is the blue - green algae Spirulina platensis that constitutes 95% of the algal population. Primary consumers vary from microscopic zooplanktons to the flamingo. The most important zooplankton is the copepod Lavenula africana though rotifers such as Brachinous dimidiatus and B. plicatilis occasionally become dominant. The lesser flamingo is the major primary consumer of Spirulina platensis while Oreochromis alcalicus grahami also grazes on it. Other important primary consumers include Hippopotamus Hippopotamus amphibius, which feed on the shoreline vegetation at night and import nutrient when they return to the open water zone.

4 - What is the Site like? (Ecological character description)

4.1 - Ecological character

Three major Rivers, the Njoro, Makalia and Nderit drain into the Lake, together with treated water from the town's sewage works and the outflows from several springs along the shore. Generally these flows cause nutrient cycling, which is of great benefit to the aquatic organisms. Diatoms and other algae that bloom occasionally form substantial amount of total biomass. The main primary producer of the open water zone is the blue - green algae Arthrospira fusiformis that constitutes 95% of the algal population. Primary consumers vary from microscopic zooplanktons to the flamingo. The most important zooplankton is the copepod Lavenula africana though rotifers such as Brachinous dimidiatus and B. plicatilis occasionally become dominant. The lesser flamingo is the major primary consumer of Arthrospira fusiformis while Oreochromis alcalicus grahami also grazes on it. Other important primary consumers include Hippopotamus Hippopotamus amphibius, which feed on the shoreline vegetation at night and import nutrient when they return to the open water zone.

The lake shores are mainly open alkaline mud with areas of sedge Cyperus laevigatus and Typha marsh around the River inflows and springs, giving way to grassland and a belt of Acacia xanthophloea woodland. The Eastern and Northern shores are characterized by a swamp of deep, soft mud kept moist by springs and seepage along the lakeside. Sedges and rushes grow here and provide shelter for a variety of animals and birds. The rushes are mostly confined to areas of relatively fresh water while the sedges tolerate more soda. Many waders especially the migrants sift through the mud for small insects. The Western and Southern shores are sandy, and are habitat for sand pipers and other water birds.

4.2 - What wetland type(s) are in the site?

Wetland types (code and name)	Local name	Ranking of extent (1: greatest - 4: least)	Area (ha) of wetland type	Justification of Criterion 1
Fresh water > Flowing water >> N: Seasonal/ intermittent/ irregular rivers/ streams/ creeks	Larmudiac river	4		
Saline, brackish or alkaline water > Lakes >> Q: Permanent saline/ brackish/ alkaline lakes	Lake Nakuru	1	5000	
Saline, brackish or alkaline water > Marshes & pools >> Sp: Permanent saline/ brackish/ alkaline marshes/ pools	Lake Nakuru marshes	3		
Fresh water > Lakes and pools >> Tp: Permanent freshwater marshes/ pools	Baharini springs	2		

Human-made wetlands

Wetland types (code and name)	Local name	Ranking of extent (1: greatest - 4: least)	Area (ha) of wetland type
8: Wastewater treatment areas	Nakuru sewarage treatment plant	3	

Other non-wetland habitats within the site	Area (ha) if known
Savannah grasslands	
Acacia xanthophloea woodlands	
Tarchonanthus camphoratus shrubland	
Euphorbia forest	

(ECD) Habitat connectivity Lake Nakuru constitute one of the extensive Kenyan Rift Lakes System under the World Heritage Sites.

4.3 - Biological components

4.3.1 - Plant species

Other noteworthy plant species

Phylum	Scientific name	Position in range / endemism / other
TRACHEOPHYTA/LILIOPSIDA	Chloris gayana	in open grassland
TRACHEOPHYTA/LILIOPSIDA	Cynodon dactylon	
TRACHEOPHYTA/LILIOPSIDA	Cyperus laevigatus	in open grassland
TRACHEOPHYTA/LILIOPSIDA	Digitaria abyssinica	in open grassland
TRACHEOPHYTA/MAGNOLIOPSIDA	Euphorbia candelabrum	
TRACHEOPHYTA/MAGNOLIOPSIDA	Olea europaea cuspidata	
TRACHEOPHYTA/MAGNOLIOPSIDA	Pluchea bequaertii	in open grassland
TRACHEOPHYTA/LILIOPSIDA	Sporobolus spicatus	in open grassland
TRACHEOPHYTA/MAGNOLIOPSIDA	Tarchonanthus camphoratus	in bushed woodlands
TRACHEOPHYTA/LILIOPSIDA	Themeda triandra	in open grassland
TRACHEOPHYTA/LILIOPSIDA	Typha domingensis	
TRACHEOPHYTA/MAGNOLIOPSIDA	Vachellia xanthophloea	
TRACHEOPHYTA/MAGNOLIOPSIDA	Vepris simplicifolia	

Invasive alien plant species

4	Impacts	Changes at RIS update
Achyranthes aspera	Actual (minor impacts)	increase
Datura stramonium	Actual (minor impacts)	No change
Lantana camara	Actual (minor impacts)	increase
Lantana trifolia	Actual (minor impacts)	increase
Lippia javanica	Actual (minor impacts)	increase
Ocimum gratissimum	Actual (minor impacts)	increase
Ocimum gratissimum gratissimum	Actual (minor impacts)	No change
Sida schimperiana	Actual (minor impacts)	increase
Solanum incanum	Actual (minor impacts)	increase
Solanum villosum	Actual (minor impacts)	No change
. Urtica massaica	Actual (minor impacts)	increase
	Datura stramonium Lantana camara Lantana trifolia Lippia javanica Ocimum gratissimum Ocimum gratissimum gratissimum Sida schimperiana Solanum incanum Solanum villosum	Actual (minor impacts) Lantana camara Actual (minor impacts) Lantana trifolia Actual (minor impacts) Lippia javanica Actual (minor impacts) Actual (minor impacts) Ocimum gratissimum Ocimum gratissimum Actual (minor impacts) Sida schimperiana Actual (minor impacts) Actual (minor impacts)

Optional text box to provide further information

Intensive habitat management strategies to reduce/manage the invasive species are being undertaken in the park. Such strategies include the mapping of invasive species distribution and manual removal of invasive species in specified management blocks. Nevertheless more funds needed to sustainably manage the invasive species throughout the year with little to no breaks between management sessions as is the case now.

4.3.2 - Animal species

Other noteworthy animal species

Phylum	Scientific name	Pop. size	Period of pop. est.	%occurrence	Position in range /endemism/other
CHORDATA/AVES	Circus macrourus				Near threatened
CHORDATA/AVES	Coracias garrulus				Near threatened
CHORDATA/AVES	Falco concolor				Near threatened
CHORDATA/AVES	Gallinago media				Near threatened
CHORDATA/AVES	Gyps africanus				Endangered
CHORDATA/AVES	Gyps rueppellii				Endangered
CHORDATA/AVES	Limosa limosa				Near threatened
CHORDATAVAVES	Oxyura maccoa				Near threatened
CHORDATAVAVES	Stephanoaetus coronatus				Near threatened
CHORDATA/AVES	Terathopius ecaudatus				Near threatened
CHORDATA/AVES	Torgos tracheliotus				Vulnerable
CHORDATA/AVES	Trigonoceps occipitalis				Vulnerable
CHORDATA/AVES	Turdus abyssinicus				Vulnerable

Invasive alien animal species

interve and animal epocies			
Phylum	Scientific name	Impacts	Changes at RIS update
CHORDATA/ACTINOPTERYGII	Oreochromis niloticus	Actual (minor impacts)	No change
CHORDATA/ACTINOPTERYGII	Oreochromis variabilis	Actual (minor impacts)	No change

4.4 - Physical components

4.4.1 - Climate

Climatic region	Subregion
A: Tropical humid climate	Aw: Tropical savanna (Winter dry season)

Due to changes in climate, Lake Nakuru rose from 40.4 km2 to 68.18 km2, a difference of 28.14 km2 or 70.28%. The rising levels of the lake coincide with the increase in rainfall in the catchment areas, with sharp increases in the surface area observed in 2014 and 2020. (GoK and UNDP 2021).

The rising water levels have also affected livelihoods through loss of grazing and farm and crop land. Reports of attacks by wild animals like snakes have also been made. Additionally, it was observed that the affected communities are at high risk of contracting waterborne diseases due to the submersion of public utilities like sewage treatment plants and other sanitation units.

4.4.2 - Geomorphic setting

a) Minimum elevation above sea level (in metres)	1756
a) Maximum elevation above sea level (in metres)	2059
	Entire river basin
	Upper part of river basin ☐
	Middle part of river basin □
	Lower part of river basin 🗹
	More than one river basin ☐
	Not in river basin
	Coastal 🗆
Please name the river basin or basins. If the s	site lies in a sub-basin, please also name the larger river basin. For a coastal/marine site, please name the sea or ocean.
ake Nakuru Catchment	

4.4.3 - Soil

)rc			

(Update) Changes at RIS update No change Increase O Decrease O Unknown O

No available information \square

Are soil types subject to change as a result of changing hydrological conditions (e.g., increased salinity or acidification)?

Please provide further information on the soil (optional)

The area surrounding Lake Nakuru catchment belongs to Tertiary - Quaternary volcanic materials associated with alkaline sediments. The main geological features found around this wetland include:

- (i) The Rift Valley bottom or lowland, in which the Lake Nakuru lies,
- (ii) The escarpments including the foot slopes and cliffs on the western side of the lake, Mau and Eburru
- (iii) The uplifted lion hill range on the eastern side and other hills,
- (iv) Menengai crater on the northern side of the lake and
- (v) Fault lines that run in a North South direction.

4.4.4 - Water regime

Water permanence

Trator pormanonos				
Presence?	Changes at RIS update			
Usually seasonal, ephemeral or intermittent water present				
Usually permanent water present	increase			

Source of water that maintains character of the site

Presence?	Predominant water source	Changes at RIS update		
Water inputs from surface water		increase		
Water inputs from groundwater		increase		
Water inputs from precipitation		increase		

Water destination

Presence?	Changes at RIS update No change	
Unknown		
Feeds groundwater	increase	

Stability of water regime

Presence?		Changes at RIS update
Water levels fluctuating (including tidal)		No change
	Water levels largely stable	increase

Please add any comments on the water regime and its determinants (if relevant). Use this box to explain sites with complex hydrology;

The lake is a closed basin experiencing rising Lake levels from 4.5 to 10 meters in the past 12 years (2010-2022)

(ECD) Connectivity of surface waters and of groundwater are done of the connectivity of surface waters and of groundwater are done of the connectivity of surface waters and of groundwater are done of the connectivity of surface waters and of groundwater are done of the connectivity of surface waters and of groundwater are done of the connectivity of surface waters and of groundwater are done of the connectivity of surface waters and of groundwater are done of the connectivity of surface waters and of groundwater are done of the connectivity of surface waters and of groundwater are done of the connectivity of surface waters and of groundwater are done of the connectivity of surface waters and of groundwater are done of the connectivity of surface waters and of groundwater are done of the connectivity of surface waters and of groundwater are done of the connectivity of surface waters and of groundwater are done of the connectivity of surface waters are

(ECD) Stratification and mixing regime The lake is stratified and there is mixing in the afternoons due to the winds

4.4.5 - Sediment regime

Sediment regime is highly variable, either seasonally or inter-annually

(Update) Changes at RIS update No change O Increase O Decrease O Unknown O

Sediment regime unknown

Please provide further information on sediment (optional):

The sedimentation is dependent on the rainfall regime in the catchment as well as the human activities

(ECD) Water turbidity and colour 27.9 and Brown colour 240C

4.4.6 - Water pH

Alkaline (pH>7.4)

Unknown \square

Please provide further information on pH (optional):

RIS for Site no. 476, Lake Nakuru, Kenya		
9.4 declined from 10.5 due to increasing discharge evidenced by rising lake levels and increased lake area		
4.4.7 - Water salinity		
Euhaline/Eusaline (30-40 g/l)		
(Update) Changes at RIS update No change □ Increase ○ Decrease ○ Unknown ○		
Unknown Unknown		
Please provide further information on salinity (optional):		
3.1 (ppt)		
(ECD) Dissolved gases in water		
7,1 mg/l		
4.4.8 - Dissolved or suspended nutrients in water Eutrophic (Update) Changes at RIS update No change Increase Decrease Unknown Unknown Unknown Unknown Decrease Provide further information on dissolved or suspended nutrients (optional):		
High nutrients levels of nitrates, phosphates and ammonia were recorded with nitrates scoring highest levels of between 3-4 mg/l		
(ECD) Water conductivity 5709.9 μs/cm		
4.4.9 - Features of the surrounding area which may affect the Site		
Please describe whether, and if so how, the landscape and ecological characteristics in the area surrounding the Ramsar Site differ from the i) broadly similar O ii) significantly different \odot site itself:		
Surrounding area has greater urbanisation or development 🗹		
Surrounding area has higher human population density 🗹		

Surrounding area has significantly different land cover or habitat types Image: Please describe other ways in which the surrounding area is different:

There is increasing urbanization, human population, settlements and land use changes driven by Nakuru City. The city is fast growing and has challenges of solid and liquid wastes which finds its way into Lake Nakuru

4.5 - Ecosystem services

4.5.1 - Ecosystem services/benefits

Provisioning Services

Ecosystem service	Examples	Importance/Extent/Significance
Genetic materials	Ornamental species (live and dead)	Medium

Surrounding area has more intensive agricultural use

Regulating Services

Ecosystem service	Examples	Importance/Extent/Significance
Maintenance of hydrological regimes	Groundwater recharge and discharge	High
Erosion protection	Soil, sediment and nutrient retention	Medium
Pollution control and detoxification	Water purification/waste treatment or dilution	Medium
Climate regulation	Regulation of greenhouse gases, temperature, precipitation and other climactic processes	Medium
Hazard reduction	Flood control, flood storage	Medium

Cultural Services

Ecosystem service	Examples	Importance/Extent/Significance
Recreation and tourism	Picnics, outings, touring	High
Recreation and tourism	Nature observation and nature-based tourism	High
Spiritual and inspirational	Cultural heritage (historical and archaeological)	Medium
Spiritual and inspirational	Aesthetic and sense of place values	Medium
Scientific and educational	Type location for a taxon	High
Scientific and educational	Long-term monitoring site	High
Scientific and educational	Major scientific study site	Medium
Scientific and educational	Important knowledge systems, importance for research (scientific reference area or site)	High
Scientific and educational	Educational activities and opportunities	High

Supporting Services

Ecosystem service	Examples	Importance/Extent/Significance
Biodiversity	Supports a variety of all life forms including plants, animals and microorganizms, the genes they contain, and the ecosystems of which they form a part	High

Other ecosystem service(s) not included above:

Cardi Goody at the Control and a solution of the Control and t
Carbon sequestration
Within the site: Residents approx. 20
•
Have studies or assessments been made of the economic valuation of ecosystem services provided by this Ramsar Site?
Where economic studies or assessments of economic valuation have been undertaken at the site, it would be helpful to provide information on where the results of such studies may be located (e.g. website links, citation of published literature):
(PDF) Environmental valuation in developing countries: The recreational value of wildlife viewing (researchgate.net)

4.5.2 - Social and cultural values

i) the site provides a model of wetland wise use, demonstrating the application of traditional knowledge and methods of management and use that maintain the ecological character of the wetland	
ii) the site has exceptional cultural traditions or records of former civilizations that have influenced the ecological character of the wetland	
iii) the ecological character of the wetland depends on its interaction with local communities or indigenous peoples	
iv) relevant non-material values such as sacred sites are present and their existence is strongly linked with the maintenance of the ecological character of the wetland	

<no data available>

4.6 - Ecological processes

<no data available>

5 - How is the Site managed? (Conservation and management)

5.1 - Land tenure and responsibilities (Managers)

5.1.1 - Land tenure/ownership

Pub	li o	OVAZI	ance	hii	_

Category	Within the Ramsar Site	In the surrounding area
National/Federal government	/	/
Local authority, municipality, (sub)district, etc.		2

Private ownership

Category	Within the Ramsar Site	In the surrounding area
Other types of private/individual owner(s)		✓

 $Provide \ further \ information \ on \ the \ land \ tenure \ / \ ownership \ regime \ (optional):$

Public land held in trust by the National Government managed by the state agency responsible for wildlife management (Kenya Wildlife Service)

5.1.2	- N	lanac	jement	auth	nority

Please list the local office / offices of any	Kenya Wildlife Service, Nakuru
agency or organization responsible for	
managing the site:	
Provide the name and/or title of the person	
people with responsibility for the wetland:	Titus Mitau, Site manager, Lake Nakuru National Park
people with responsibility for the wettand.	
Postal address:	P.O Box 535-20100 Nakuru
Postal address.	
E-mail address:	swlakenakuru@kws.go.ke

5.2 - Ecological character threats and responses (Management)

5.2.1 - Factors (actual or likely) adversely affecting the Site's ecological character

Human settlements (non agricultural)

Factors adversely affecting site	Actual threat	Potential threat	Within the site	Changes	In the surrounding area	Changes
Housing and urban areas	High impact	High impact		No change	2	increase
Commercial and industrial areas	High impact	High impact		No change	2	increase
Tourism and recreation areas	Low impact	Medium impact	/	increase	2	increase
Unspecified development	High impact	High impact		No change	2	No change

Water regulation

vator regulation						
Factors adversely affecting site	Actual threat	Potential threat	Within the site	Changes	In the surrounding area	Changes
Drainage	High impact	High impact	✓	increase	✓	increase
Water abstraction	Medium impact	Medium impact		No change	✓	increase

Agriculture and aquaculture

Factors adversely affecting site	Actual threat	Potential threat	Within the site	Changes	In the surrounding area	Changes
Annual and perennial non-timber crops	High impact	High impact		No change	/	increase
Wood and pulp plantations	High impact	High impact		No change	2	increase
Livestock farming and ranching			/		/	

energy production and mining							
Factors adversely affecting site	Actual threat	Potential threat	Within the site	Changes	In the surrounding area	Changes	
Mining and quarrying	High impact	High impact		No change	✓	No change	
Renewable energy	Low impact	Medium impact		No change	✓	increase	

Transportation and service corridors

Factors adversely affecting site	Actual threat	Potential threat	Within the site	Changes	In the surrounding area	Changes
Roads and railroads	Low impact	High impact	✓	increase	✓	increase
Utility and service lines (e.g., pipelines)	Low impact	High impact	2	No change	2	increase

Biological resource use

Factors adversely affecting site	Actual threat	Potential threat	Within the site	Changes	In the surrounding area	Changes
Hunting and collecting terrestrial animals	Low impact	Medium impact		No change	2	increase
Gathering terrestrial plants					/	
Logging and wood harvesting					/	
Fishing and harvesting aquatic resources	Low impact	Low impact	✓	No change	2	increase

Human intrusions and disturbance

Factors adversely affecting site	Actual threat	Potential threat	Within the site	Changes	In the surrounding area	Changes
Recreational and tourism activities	Low impact	High impact	/	No change	/	increase

Natural system modifications

Natural system modifications						
Factors adversely affecting site	Actual threat	Potential threat	Within the site	Changes	In the surrounding area	Changes
Fire and fire suppression	High impact	High impact	✓	No change	✓	No change
Dams and water management/use	Low impact	Medium impact	/	No change	/	increase
Vegetation clearance/ land conversion	Medium impact	Medium impact		No change	>	increase
Unspecified/others			/		✓	

Invasive and other problematic species and genes

Factors adversely affecting site	Actual threat	Potential threat	Within the site	Changes	In the surrounding area	Changes
Invasive non-native/ alien species	High impact	High impact	2	increase	✓	increase
Problematic native species	Low impact	Medium impact	2	No change	✓	increase

Pollution

Factors adversely affecting site	Actual threat	Potential threat	Within the site	Changes	In the surrounding area	Changes
Household sewage, urban waste water	High impact	High impact	✓	increase	✓	increase
Industrial and military effluents	High impact	High impact	/	increase	✓	increase
Agricultural and forestry effluents	High impact	High impact	✓	increase	✓	increase
Garbage and solid waste	High impact	High impact	/	increase	✓	increase
Air-borne pollutants	High impact	High impact	V	No change	✓	No change

Climate change and severe weather

Factors adversely affecting site	Actual threat	Potential threat	Within the site	Changes	In the surrounding area	Changes
Habitat shifting and alteration	High impact	High impact	/	decrease	/	increase
Droughts	High impact	High impact	✓	No change	✓	No change
Storms and flooding	High impact	High impact	✓	increase	✓	increase

5.2.2 - Legal conservation status

Global legal designations

Designation type	Name of area	Online information url	Overlap with Ramsar Site
Other global designation	Lake Nakuru Important Bird Area	https://www.datazone.birdlife.or g/userfiles/files/lBAs/monitorin gPDFs/2011-Kenya	whole
Other global designation	Lake Nakuru Ramsar Site	htttps://www.ramsar.org/wetland/ kenya	whole
World Heritage site	Kenya lake system in the Great Rift Valley	https://whc.unesco.org/en/list/1 060	whole
World Heritage site	Lake Nakuru National Park	www.whc.unesco.org/en/list/1060	whole

National legal designations

Designation type	Name of area	Online information url	Overlap with Ramsar Site
Bird Sanctuary			whole
Forestry Conservation Areas			whole
National Park	Lake Nakuru National Park	www.kws.go.ke	whole

Non-statutory designations

Designation type	Name of area	Online information url	Overlap with Ramsar Site
Important Bird Area	Lake Nakuru National Park	www.birdlife.org	whole

5.2.3 - IUCN protected areas categories (2008)

	la Strict Nature Reserve
	Ib Wilderness Area: protected area managed mainly for wilderness protection
ø	II National Park: protected area managed mainly for ecosystem protection and recreation
	III Natural Monument: protected area managed mainly for conservation of specific natural features
	IV Habitat/Species Management Area: protected area managed mainly for conservation through management intervention
	V Protected Landscape/Seascape: protected area managed mainly for landscape/seascape conservation and recreation
	VI Managed Resource Protected Area: protected area managed mainly for the sustainable use of natural ecosystems

5.2.4 - Key conservation measures

Legal protection

9 F	
Measures	Status
Legal protection	Implemented

Habitat

Measures	Status
Improvement of water quality	Partially implemented
Habitat manipulation/enhancement	Partially implemented
Hydrology management/restoration	Partially implemented
Catchment management initiatives/controls	Partially implemented
Soil management	Partially implemented
Re-vegetation	Partially implemented

Species

Species	
Measures	Status
Threatened/rare species management programmes	Implemented
Control of invasive alien plants	Partially implemented
Reintroductions	Implemented

Human Activities

Measures	Status
Communication, education, and participation and awareness activities	Implemented
Research	Implemented
Management of water abstraction/takes	Partially implemented
Regulation/management of wastes	Partially implemented
Regulation/management of recreational activities	Implemented
Harvest controls/poaching enforcement	Implemented
Livestock management/exclusion (excluding fisheries)	Implemented

5.2.5 - Management planning

Is there a site-specific management plan for the site? Yes

Has a management effectiveness assessment been undertaken for the site?

If the site is a formal transboundary site as indicated in section Data and location > Site location, are there shared management planning Yes O No opposes see with another Contracting Party?

Please indicate if a Ramsar centre, other educational or visitor facility, or an educational or visitor programme is associated with the site:

Research facilities

- (i) Research Station Lake Nakuru National Park has a full fledged research station.
- (ii) Water Quality Monitoring Laboratory A modern laboratory is in place that can monitor potable water, effluents; sewage works efficiency, lake and river waters.

The Park has two education centers for school visits and tailor - made training.

Visitor facilities in the site: 2 Lodges with a total occupancy capacity of 240 beds, picnic sites, special and public campsites and view points.

5.2.6 - Planning for restoration

Is there a site-specific restoration plan? Yes, there is a plan $\,$

5.2.7 - Monitoring implemented or proposed

Monitoring	Status
Water regime monitoring	Implemented
Water quality	Implemented
Plant community	Implemented
Animal community	Implemented
Animal species (please specify)	Implemented
Birds	Implemented

Rhino surveillance done on a 24 - hour basis. Sighting and data on habitat utilization, social characteristics and feeding patterns are recorded. Individual Animal location and movement patterns are mapped using GIS.

Meteorology: Meteorological variables currently being monitored are rainfall, temperature, relative humidity, evaporation, radiation and wind speed.

6 - Additional material

6.1 - Additional reports and documents

6.1.1 - Bibliographical references

- 1.Government of Kenya and UNDP (2021) Rising Water Levels in Kenya's Rift Valley Lakes, Turkwel Gorge Dam and Lake Victoria
- 2. Ireeene M. et.al (2020) January 2020 Waterbird Count Results in the RiftValley, Nairobi, Central, Eastern, and Coastal Kenya
- 3. Kiogora T. et.al (2020) Assessment of the fish species and water quality in lake nakuru: cotoxicological study of the potential role of heavy metals in the recent fish die-offs

6.1.2 - Additional reports and documents

i. taxonomic lists of plant and animal species occurring in the site (see section 4.3)

<1 file(s) uploaded:

ii. a detailed Ecological Character Description (ECD) (in a national format)

iii. a description of the site in a national or regional wetland inventory

iv. relevant Article 3.2 reports

v. site management plan

vi. other published literature

<2 file(s) uploaded>

6.1.3 - Photograph(s) of the Site

Please provide at least one photograph of the site:



aerial v iew Lake nakuru showing submerged roads after lake water lev el rise (Martha Mutiso, marthnzisa@gmail.com, 07-08-2013)



the background (Martha Mutiso, marthnzisa@gmail.com, 09-10-2006)



Aerial view of Lake Nakuru (Martha Mutiso, marthnzisa@gmail.com, 09-10-2006)



Flamingos in Lake Nakuru in the background (Martha Mutiso, marthnzisa@gmail.com, 09-10-2006)

6.1.4 - Designation letter and related data

Designation letter

<1 file(s) uploaded>

Date of Designation 1990-06-05