

# Ramsar Information Sheet

Published on 8 May 2023 Update version, previously published on : 8 March 2018

# Norway Kvisleflået



Designation date 6 August 2002 Site number 1193

Coordinates 61°48'25"N 12°06'11"E

Area 5 682,00 ha

### Color codes

Fields back-shaded in light blue relate to data and information required only for RIS updates.

Note that some fields concerning aspects of Part 3, the Ecological Character Description of the RIS (tinted in purple), are not expected to be completed as part of a standard RIS, but are included for completeness so as to provide the requested consistency between the RIS and the format of a 'full' Ecological Character Description, as adopted in Resolution X.15 (2008). If a Contracting Party does have information available that is relevant to these fields (for example from a national format Ecological Character Description) it may, if it wishes to, include information in these additional fields.

### 1 - Summary

#### Summary

The site is a large, flat mire area bordering Sweden in Engerdal. Several large and flat mire expanses occur, although it is more typical with poorer small mires and low pine-covered knolls. Birch Betula grows at the edge of the mires as well as along rivers and streams. The area is dominated by numerous small pools and ponds. Within the mire, a series of moraine ridges run south-east to north-west. In the southern part, we find old forest of mainly spruce and some birch.

Kviseflået is an important area for wetland birds, with 20-30 breeding species recorded. The mosaic landscape is ideal for the greenshank Tringa nebularis and the whimbrel Numenius phaeopus. After the ice melts in the spring, several duck species stage on route to upland breeding sites around the reserve. Several regionally uncommon species breed, or are suspected to have bred, in the reserve including the whooper swan Cygnus cygnus, the bean goose Anser fabalis, the broad-billed sandpiper Limicola falcinellus and the jack snipe Lymnocryptes minimus.

### 2 - Data & location

## 2.1 - Formal data

2.1.1 - Name and address of the com	piler of this RIS
Responsible compiler	
Institution/agency	Norwegian Envirnonment Agency
Postal address	P.O. Box 5672 Torgarden, N-7485 Trondheim, Norway
National Ramsar Administrati	ve Authority
Postal address	Postboks 5672 Sluppen Trondheim Norway
2.1.2 - Period of collection of data and	d information used to compile the RIS
From year	2004
To year	2021
2.1.3 - Name of the Ramsar Site	
Official name (in English, French or Spanish)	Kvisleflået
(Update) A	d area of the Site since its designation or earlier update  Changes to Site boundary Yes O No   B. Changes to Site area No change to area
(Update) For secretariat only: T	
2.1.5 - Changes to the ecological cha (Update) 6b i. Has the ecological character of tapplicable Criteria) change	he Ramsar Site (including
2.2 - Site location	
2.2.1 - Defining the Site boundaries	
b) Digital map/image	
<1 file(s) uploaded>	
Former maps	0
Boundaries description  The Ramsar Site is equal with the boundaries description	order of the Kvisleflået Nature Reserve.
2.2.2 - General location	
a) In which large administrative region does the site lie?	Innlandet
b) What is the nearest town or population centre?	Engerdal
2.2.3 - For wetlands on national bound	daries only
a) Does the wetland extend onto the ter	rritory of one or more other countries? Yes  No O
b) Is the site adjacent to another design territory of a	gnated Ramsar Site on the Yes O No O another Contracting Party?

Official area, in hectares (ha): 5682

2.2.4 - Area of the Site

Area, in hectares (ha) as calculated from 5683.655 GIS boundaries

### 2.2.5 - Biogeography

### Biogeographic regions

Regionalisation scheme(s)	Biogeographic region
EU biogeographic regionalization	1. Boreal
Other scheme (provide name below)	Northern boreal vegetation zone, transitional zone (Nb-OC)

### Other biogeographic regionalisation scheme

- 1. Biogeographical regions of Europe, European Environment Agency, 2005
- 2. Zonal division showing the variation in vegetation from south to north and from the lowlands to the mountains, and sectional graduation showing the variation between the coast and inland (In: Moen, A. 1998. Nasjonalatlas for Norge; vegetasjon. Statens kartverk, Hønefoss).

## 3 - Why is the Site important?

### 3.1 - Ramsar Criteria and their justification

☑ Criterion 1: Representative, rare	or unique natural or near-natural wetland types			
Other ecosystem services provided	Extensive mires like this function as important carbon storages.			
	A large mire complex with a large number of small pools and ponds, typical and representative for the			
Other reasons	continental part of southern Norway.			
☑ Criterion 2 : Rare species and the	reatened ecological communities			
Critorian 3 : Biological diversity				

The site is one of few breeding sites for the jack snipe Lymnocryptes minimus in southern Norway. In Justification addition several easterly and north-easterly species nest such as the whooper swan Cygnus cygnus, the bean goose Anser fabalis, the Siberian tit Parus cinctus and the rustic Bunting Emberiza rustica.

☑ Criterion 4 : Support during critical life cycle stage or in adverse conditions

3.2 - Plant species whose presence relates to the international importance of the site

Phylum	Scientific name	Criterion 2	Criterion 3	Criterion 4	Red CITES Appendix I	Other status	Justification
Fungi							
BASIDIOMYCOTA/ AGARICOMYCETES	Amylocystis lapponicus	V				National Red List: Considered as VU	
ASCOMYCOTA / NOT ASSIGNED	Chaenotheca laevigata	V				National Red List: Considered as VU	
ASCOMYCOTA / NOT ASSIGNED	Chaenotheca phaeocephala	<b></b> ✓				National Red List: Considered as VU	
BASIDIOMYCOTA/ AGARICOMYCETES	Crustoderma dryinum	V				National Red List: Considered as VU	
ASCOMYCOTA / LECANOROMYCETES	Cyphelium inquinans	V				National Red List: Considered as VU	
ASCOMYCOTA / LECANOROMYCETES	Cyphelium karelicum	<b></b> ✓				National Red List: Considered as VU	
BASIDIOMYCOTA / AGARICOMYCETES	Diplomitoporus crustulinus	V				National Red List: Considered as VU	
BASIDIOMYCOTA / AGARICOMYCETES	Gloeophyllum protractum	V				National Red List: Considered as VU	
BASIDIOMYCOTA/ AGARICOMYCETES	Laurilia sulcata	<b></b> ✓				National Red List: Considered as VU	
BASIDIOMYCOTA / AGARICOMYCETES	Perenniporia subacida	V				National Red List: Considered as EN	
BASIDIOMYCOTA/ AGARICOMYCETES	Skeletocutis chrysella	$\checkmark$				National Red List: Considered as VU	
BASIDIOMYCOTA/ AGARICOMYCETES	Skeletocutis odora	<b>/</b>				National Red List: Considered as VU	
BASIDIOMYCOTA/ AGARICOMYCETES	Skeletocutis stellae	<b></b> ✓				National Red List: Considered as VU	

Protection status is given according to the National Red List 2021.	

3.3 - Animal species whose presence relates to the international importance of the site

3.3 - Ani	mai species			ese			ates	to the international im	ipori	tance o	t the site	<b>e</b>	
Phylum	Scientific name	Spec qualifies crite 2 4	unde rion	un	der c	butes riterio	Size	Period of pop. Est. % occurrence	IUCN Red List	CITES Appendix I	CMS Appendix I	Other Status	Justification
Others													
CHORDATA / MAMMALIA	Gulo gulo						]		LC			National Red List: Considered as EN	
CHORDATA / MAMMALIA	Lynx lynx						]		LC			National Red List: Considered as EN	
CHORDATA / MAMMALIA	Ursus arctos						]		LC	V		National Red List: Considered as EN	
Birds													
CHORDATA/ AVES	Anas crecca						)		LC				Criterion 4: This species breeds at the site.
CHORDATA/ AVES	Anser fabalis			Z			)		LC			National Red List: Considered as EN	Criterion 3: This species is observed from time to time in the area.
CHORDATA / AVES	Aythya fuligula						]		LC				Criterion 4: This species breeds at the site.
CHORDATA / AVES	Bucephala clangula						]		LC				Criterion 4: This species breeds at the site.
CHORDATA / AVES	Circus cyaneus	<b>V</b>		<b>V</b>			)		LC			National Red List: Considered as EN	Criterion 4: This species is recorded during the breeding season and may breed occasionally.
CHORDATA/ AVES	Cygnus cygnus	<b>V</b>		V			)		LC			Annex II, Bern Convention	Criterion 3 & 4: This species breeds here. Nesting by Whooper Swan Cygnus cygnus in 1991 was the first record in the county.
CHORDATA/ AVES	Gavia arctica						]		LC				Criterion 4: This species breeds here.
CHORDATA / AVES	Grus grus			V			)		LC				Criterion 4: This species nests here.
CHORDATA/ AVES	Limicola falcinellus	1		V			]		LC			Annex II, Bern Convention	Criterion 4: This species nests here.
CHORDATA/ AVES	Lymnocryptes minimus			V			)		LC				Criterion 3 & 4: The species is regularly observed, and probably breeds on the site.
CHORDATA /	Numenius phaeopus						)		LC				Criterion 4: This species breeds at the site.
CHORDATA/ AVES	Philomachus pugnax	<b>V</b>		V			]		LC			National Red List: Considered as VU	Criterion 4: This species is recorded during the breeding season and may breed occasionally.
CHORDATA/ AVES	Poecile cinctus			<b>V</b>			1		LC				Criterion 3 & 4: This species nests here. Western limit of its range.
CHORDATA / AVES	Tringa glareola	<b>V</b>		V			)		LC			Annex II, Bern Convention	Criterion 4: Common breeder on the site.
CHORDATA / AVES	Tringa nebularia						)		LC				Criterion 4: This species breeds at the site.
CHORDATA / AVES	Vanellus vanellus	<b>V</b>		V			)		NT			National Red List: Considered as CR	Crtiterion 4: This species breeds on the site.

1) Percentage of the total biogeographic population at the site

Protection status is given according to the National Red List 2021.

### 3.4 - Ecological communities whose presence relates to the international importance of the site

<no data available>

### 4 - What is the Site like? (Ecological character description)

### 4.1 - Ecological character

Characteristic for the area is a mosaic of small nutrient-poor mires, and low pine-clad knolls, as well as large open mire areas such as at Storkjølen and Rådløskjølen. The vegetation varies from nutrient-poor mire/intermediate mire vegetation, low pine woods on the moraine ridges and willow Salix- swamp woodland with some birch Betula along rivers and edges of mires. A substantial part of the forest is old forest/natural forest, With a very high diversity of nationally Red-Listed fungi and lichen species connected to this forest type. In places there are areas of wet mixed woodland with characteristics resembling ancient woodland. The many pools in the area are important for birdlife, although they have little aquatic vegetation. The birdlife includes a good representation of breeding waterbirds - albeit in small numbers.

### 4.2 - What wetland type(s) are in the site?

Inland wetlands

Wetland types (code and name)	Local name	Ranking of extent (1: greatest - 4: least)	Area (ha) of wetland type	Justification of Criterion 1
Fresh water > Flowing water >> L: Permanent inland deltas		0		
Fresh water > Flowing water >> M: Permanent rivers/ streams/ creeks				
Fresh water > Lakes and pools  >> O: Permanent freshwater lakes		4		
Fresh water > Lakes and pools >> Tp: Permanent freshwater marshes/ pools		3		Representative
Fresh water > Lakes and pools >> Ts: Seasonal/ intermittent freshwater marshes/ pools on inorganic soils		0		
Fresh water > Marshes on peat soils >> U: Permanent Non- forested peatlands		1		Representative
Fresh water > Marshes on inorganic soils >> Xf: Freshwater, tree-dominated wetlands		2		
Fresh water > Marshes on peat soils >> Xp: Permanent Forested peatlands		0		

### 4.3 - Biological components

### 4.3.1 - Plant species

Other noteworthy plant species

Phylum	Scientific name	Position in range / endemism / other
ASCOMYCOTALECANOROMYCETES	Alectoria sarmentosa	National Red List: Considered as NT
BASIDIOMYCOTA/AGARICOMYCETES	Antrodia albobrunnea	National Red List: Considered as NT
BASIDIOMYCOTA/AGARICOMYCETES	Antrodia xantha	National Red List: Considered as LC
ASCOMYCOTALECANOROMYCETES	Bryoria nadvornikiana	National Red List: Considered as NT
ASCOMYCOTANOT ASSIGNED	Chaenotheca subroscida	National Red List: Considered as NT
BASIDIOMYCOTA/AGARICOMYCETES	Chaetoderma luna	National Red List: Considered as NT
BASIDIOMYCOTA/AGARICOMYCETES	Chaetodermella luna	National Red List: Considered as NT
ASCOMYCOTALECANOROMYCETES	Cladonia parasitica	National Red List: Considered as NT
BASIDIOMYCOTA/AGARICOMYCETES	Crustoderma corneum	National Red List: Considered as VU
BASIDIOMYCOTA/AGARICOMYCETES	Cystostereum murrayi	National Red List: Considered as LC
BASIDIOMYCOTA/AGARICOMYCETES	Fomitopsis rosea	National Red List: Considered as NT
TRACHEOPHYTA/LILIOPSIDA	Gymnadenia conopsea	National Red List: Considered as LC
ASCOMYCOTALECANOROMYCETES	Hypogymnia bitteri	National Red List: Considered as NT
BASIDIOMYCOTA/AGARICOMYCETES	Inonotus Ieporinus	National Red List: Considered as NT
ASCOMYCOTALECANOROMYCETES	Letharia vulpina	National Red List: Considered as NT
ASCOMYCOTA/NOT ASSIGNED	Microcalicium ahlneri	National Red List: Considered as NT
BASIDIOMYCOTA/AGARICOMYCETES	Odonticium romellii	National Red List: Considered as NT
BASIDIOMYCOTA/AGARICOMYCETES	Phlebia mellea	National Red List: Considered as NT
ASCOMYCOTA/NOT ASSIGNED	Pseudographis pinicola	National Red List: Considered as LC
BASIDIOMYCOTA/AGARICOMYCETES	Pseudomerulius aureus	National Red List: Considered as NT
BASIDIOMYCOTA/AGARICOMYCETES	Skeletocutis borealis	National Red List: Considere as DD
BASIDIOMYCOTA/AGARICOMYCETES	Trichaptum laricinum	National Red List: Considered as NT

### 4.3.2 - Animal species

Other noteworthy animal species

Phylum	Scientific name	Pop. size	Period of pop. est.	% occurrence	Position in range /endemism/other
CHORDATA/AVES	Cuculus canorus				Possibly breeding at the site. National Red List Status: NT
CHORDATAVAVES	Pluvialis apricaria				Possibly breeding on the site.
CHORDATA/AVES	Tringa totanus				Possibly breeding on the site.

### 4.4 - Physical components

### 4.4.1 - Climate

Climatic region	Subregion
D: Moist Mid-Latitude climate with cold winters	Dfc: Subarctic (Severe winter, no dry season, cool summer)

Mineral (Update) Changes at RIS update No change Increase Decrease Unknown O Organic (Update) Changes at RIS update No change Increase Decrease Unknown O No available information  Are soil types subject to change as a result of changing hydrological conditions (e.g., increased salinity or acidification)?  Please provide further information on the soil (optional)  Peat soils dominate in the large areas of mires, although the moraine ridges with pine Pinus sylvestris woodland contain mineral soils.	The climate is slightly winters.	continental with rather a lot c	of precipitation (ca. 700 mm p.a.) and relatively warm, but short, summers and extremely cold
a) Maximum elevation above sea level (in metres)    To 8	4 4 2 - Geomorphic set	ting	
a) Maximum elevation above sea level (in metres)   1000	·		
Entire river basin   Upper part of river basin   Upper part of river basin   Upper part of river basin   Lower part of river basin   Lower part of river basin   Not in river basin   Not in river basin   Not in river basin   Coastal   Please name the river basin or basins. If the site lies in a sub-basin, please also name the larger river basin. For a coastal/marine site, please name the sea or ocean.  The mire belongs to the catchement area Datelven, and it drains into the river Datelven in Sweden.  I.4.3 - Soil  Mineral (Update) Changes at RIS update No change (Increase Conditions) Changes at RIS update No change increase Conditions (e.g., increased salinity or addification)?  Are soil types subject to change as a result of changing hydrological conditions (e.g., increased salinity or addification)?  Please provide further information on the soil (optional)  Peat soils dominate in the large areas of mires, although the moraine ridges with pine Pinus sylvestris woodland contain mineral soils.  4.4.4 - Water regime  Vater permanence  Presence? Changes at RIS update  Usually permanent water	a) willillillillilli elevation at	1 /58	
Upper part of river basin   Middle part of river basin   Lower part of river basin   Not in river basin   Not in river basin   Coastal   Please name the river basin or basins. If the site lies in a sub-basin, please also name the larger river basin. For a coastal/marine site, please name the sea or ocean.  The mire belongs to the catchement area Dalelven, and it drains into the river Dalelven in Sweden.  Mineral   (Update) Changes at RIS update No change  Increase  Decrease  Unknown   Organic    (Update) Changes at RIS update No change  Increase  Decrease  Unknown   No available information Are soil types subject to change as a result of changing hydrological conditions (e.g., increased salinity or addification)?  Please provide further information on the soil (optional)  Peat soils dominate in the large areas of mires, although the moraine ridges with pine Pinus sylvestris woodland contain mineral soils.	a) Maximum elevation ab	11000	
Middle part of river basin   Lower part of river basin   Not in river ba		Entire riv	wer basin □
Lower part of river basin   More than one river basin   Not in river basin   Not in river basin   Not in river basin   Coastal   Please name the river basin or basins. If the site lies in a sub-basin, please also name the larger river basin. For a coastal/marine site, please name the sea or ocean.  The mire belongs to the catchement area Dalelven, and it drains into the river Dalelven in Sweden.    1.4.3 - Soil		Upper part of riv	ver basin ☑
More than one river basin    Not in river basin   Coastal		Middle part of riv	ver basin 🗆
Not in river basin  Coastal  Please name the river basin or basins. If the site lies in a sub-basin, please also name the larger river basin. For a coastal/marine site, please name the sea or ocean.  The mire belongs to the catchement area Dalelven, and it drains into the river Dalelven in Sweden.    1.4.3 - Soil		Lower part of riv	ver basin 🗆
Please name the river basin or basins. If the site lies in a sub-basin, please also name the larger river basin. For a coastal/marine site, please name the sea or ocean.  The mire belongs to the catchement area Dalelven, and it drains into the river Dalelven in Sweden.    1.4.3 - Soil		More than one riv	ver basin 🗆
Please name the river basin or basins. If the site lies in a sub-basin, please also name the larger river basin. For a coastal/marine site, please name the sea or ocean.  The mire belongs to the catchement area Dalelven, and it drains into the river Dalelven in Sweden.    1.4.3 - Soil		Not in riv	ver basin 🗆
The mire belongs to the catchement area Dalelven, and it drains into the river Dalelven in Sweden.    A.3 - Soil			Coastal
Mineral (Update) Changes at RIS update No change Increase Decrease Unknown Organic (Update) Changes at RIS update No change Increase Decrease Unknown Organic (Update) Changes at RIS update No change Increase Decrease Unknown Organic Decrease Unknown Organic Decrease Organic Dec	Please name the river basin	or basins. If the site lies in a sub-b	asin, please also name the larger river basin. For a coastal/marine site, please name the sea or ocean.
Mineral (Update) Changes at RIS update No change Increase Decrease Unknown Organic (Update) Changes at RIS update No change Increase Decrease Unknown Organic (Update) Changes at RIS update No change Increase Decrease Unknown Organic No available information Are soil types subject to change as a result of changing hydrological conditions (e.g., increased salinity or acidification)? Yes No Please provide further information on the soil (optional)  Peat soils dominate in the large areas of mires, although the moraine ridges with pine Pinus sylvestris woodland contain mineral soils.	The mire belongs to the	e catchement area Dalelver	n, and it drains into the river Dalelven in Sweden.
Organic (Update) Changes at RIS update No change Increase Decrease Unknown No available information  No available information  Are soil types subject to change as a result of changing hydrological conditions (e.g., increased salinity or acidification)? Yes No Please provide further information on the soil (optional)  Peat soils dominate in the large areas of mires, although the moraine ridges with pine Pinus sylvestris woodland contain mineral soils.  4.4.4 - Water regime  Water permanence  Presence? Changes at RIS update  Usually permanent water	4.4.3 - Soil		Mineral ☑
(Update) Changes at RIS update  No change ● Increase ○ Decrease ○ Unknown ○  No available information  Are soil types subject to change as a result of changing hydrological conditions (e.g., increased salinity or acidification)?  Please provide further information on the soil (optional)  Peat soils dominate in the large areas of mires, although the moraine ridges with pine Pinus sylvestris woodland contain mineral soils.  4.4.4 - Water regime  Water permanence  Presence? Changes at RIS update  Usually permanent water		(Update) Changes at RI	Supdate No change  Increase  ODecrease  OUnknown  O
Are soil types subject to change as a result of changing hydrological conditions (e.g., increased salinity or acidification)?  Please provide further information on the soil (optional)  Peat soils dominate in the large areas of mires, although the moraine ridges with pine Pinus sylvestris woodland contain mineral soils.  4.4.4 - Water regime  Vater permanence  Presence? Changes at RIS update  Usually permanent water			Organic ☑
Are soil types subject to change as a result of changing hydrological conditions (e.g., increased salinity or acidification)?  Please provide further information on the soil (optional)  Peat soils dominate in the large areas of mires, although the moraine ridges with pine Pinus sylvestris woodland contain mineral soils.  4.4.4 - Water regime  Water permanence  Presence? Changes at RIS update  Usually permanent water		(Update) Changes at RI	Supdate No change
Please provide further information on the soil (optional)  Peat soils dominate in the large areas of mires, although the moraine ridges with pine Pinus sylvestris woodland contain mineral soils.  4.4.4 - Water regime  Water permanence  Presence? Changes at RIS update  Usually permanent water		No available info	ormation
Peat soils dominate in the large areas of mires, although the moraine ridges with pine Pinus sylvestris woodland contain mineral soils.  1.4.4 - Water regime  Water permanence  Presence? Changes at RIS update  Usually permanent water	Are soil types subject to condition	change as a result of changing hyd	rological Yes O No
1.4.4 - Water regime  Vater permanence Presence? Changes at RIS update  Usually permanent water	Please provide further inform	nation on the soil (optional)	
Vater permanence Presence? Changes at RIS update Usually permanent water	Peat soils dominate in	the large areas of mires, al	though the moraine ridges with pine Pinus sylvestris woodland contain mineral soils.
Usually permanent water	4.4.4 - Water regime Water permanence	Changes at DIS undate	
present	Usually permanent water	onanges at the update	
	present		

Presence?	Changes at RIS update
Usually permanent water present	
prosent	

obaros or mater anat mannamo oriaraster or are one			
	Presence?	Predominant water source	Changes at RIS update
	Water inputs from surface water		No change
	Water inputs from precipitation		No change

### Stability of water regime

Presence?	Changes at RIS update	
Water levels largely stable	No change	

Please add any comments on the water regime and its determinants (if relevant). Use this box to explain sites with complex hydrology.

The water levels in the mire areas is rather permanent and stable.

Large mires such as this are important water reservoirs. They provide stability in water drainage in the watercourse by acting as reservoirs in drought periods and as flood barriers during snow melt and periods of heavy precipitation.

### 4.4.5 - Sediment regime

### Other ecosystem service(s) not included above:

Large mires such as this are important water reservoirs. They provide stability in water drainage in the watercourse by acting as reservoirs in drought periods and as flood barriers during snow melt and periods of heavy precipitation.

Some berry picking (in particular cloudberry) takes place, as well as hunting and fishing.

There is limited hobby fishing and hunting within the area. Domestic livestock graze the area - mainly sheep although also some cattle.

Have studies or assessments been made of the economic valuation of Yes O No O Unknown 

One Of the economic valuation of Yes O No O Unknown ecosystem services provided by this Ramsar Site?

i) the site provides a model of wetland wise use, demonstrating the application of traditional knowledge and methods of management and use that maintain the ecological character of the wetland	
ii) the site has exceptional cultural traditions or records of forme civilizations that have influenced the ecological character of the wetland	
iii) the ecological character of the wetland depends on its interaction with local communities or indigenous people:	_
iv) relevant non-material values such as sacred sites are present and their existence is strongly linked with the maintenance of the ecological character of the wetland	

<no data available>

# 4.6 - Ecological processes

<no data available>

### 5 - How is the Site managed? (Conservation and management)

### 5.1 - Land tenure and responsibilities (Managers)

#### 5.1.1 - Land tenure/ownership

Private ownership

Category		Within the Ramsar Site	In the surrounding area	
	Other types of private/individual owner(s)	✓	✓	

Provide further information on the land tenure / ownership regime (optional).

within the Ramsar site: Private in the surrounding area: Private

### 5.1.2 - Management authority

Please list the local office / offices of any County governor of Innlandet agency or organization responsible for managing the site:

Statsforvalteren i Innlandet

Postal address: Pb. 987

N-2604 LILLEHAMMER

E-mail address: sfinpost@statsforvalteren.no

### 5.2 - Ecological character threats and responses (Management)

### 5.2.1 - Factors (actual or likely) adversely affecting the Site's ecological character

Please describe any other threats (optional):

within the Ramsar site:

None known. Grazing intensity from livestock is low and has little or no impact on the vegetation.

in the surrounding area: None known.

### 5.2.2 - Legal conservation status

National legal designations

Designation type	Name of area	Online information url	Overlap with Ramsar Site
Nature Reserve	Kviseflået		whole

### 5.2.3 - IUCN protected areas categories (2008)

la Strict Nature Reserve

lb Wilderness Area: protected area managed mainly for wilderness protection

II National Park: protected area managed mainly for ecosystem  $\hfill\Box$ protection and recreation

III Natural Monument: protected area managed mainly for conservation of specific natural features

IV Habitat/Species Management Area: protected area managed mainly  $\hfill\Box$ for conservation through management intervention

V Protected Landscape/Seascape: protected area managed mainly for landscape/seascape conservation and recreation

VI Managed Resource Protected Area: protected area managed mainly for the sustainable use of natural ecosystems

### 5.2.4 - Key conservation measures

Legal protection		
Measures	Status	
Legal protection	Implemented	

#### Other

The reserve was expanded in 2005 (increased to 5682 ha).

Conservation measures proposed but not yet implemented:

The area borders a similar wetland area (Ramsar site) on the Swedish side of the border. There have been some initial efforts to coordinate the management of the transfrontier conservation area.

### 5.2.5 - Management planning

Is there a site-specific management plan for the site? In preparation

Has a management effectiveness assessment been undertaken for the site?

If the site is a formal transboundary site as indicated in section Data and location > Site location, are there shared management planning Yes O No 

processes with another Contracting Party?

### 5.2.6 - Planning for restoration

Is there a site-specific restoration plan? No need identified

### 5.2.7 - Monitoring implemented or proposed

<no data available>

### 6 - Additional material

### 6.1 - Additional reports and documents

#### 6.1.1 - Bibliographical references

Artsdatabanken (2021, 24. november). Norsk rødliste for arter 2021. https://www.artsdatabanken.no/lister/rodlisteforarter/2021

#### Botanical and management plans:

Myhr, S. 1973. Kvisleflået. Inventeringer i forbindelse med Landsplanen for verneverdige områder/forekomster. Miljøverndepartementet, 13 pp. (In Norwegian - on national plan for important conservation areas).

#### Birds:

Bekken, J. 1987. Ornitologiske registreringer i 11 våtmarksreservater 1985-86. Fylkesmannen i Hedmark, Miljøvernavd. Rapport nr. 13: 1-43. (In Norwegian – bird recording in 11 wetland reserves in Hedmark county).

Bekken, J. 2001. Fugler og pattedyr i 18 våtmarksreservater i Hedmark. Fylkesmannen i Hedmark, Miljøvernavd. Rapport nr. 8/2001: 1-122. (In Norwegian – bird and mammal recording in 18 wetland reserves in Hedmark county).

Bekken, J. 2013. Fugler i 20 våtmarksreservater i Hedmark 2000-2012. Fylkesmannen i Hedmark, Miljøvernavd. Rapport nr. 2/2013: 1-125. (In Norwegian - bird registrations in 20 Wetland reserves in Hedmark County)

#### Geology:

Sollid, J. L. & Kristiansen, K. 1982. Hedmark fylke. Kvartærgeologisk verneverdige områder. Universitetet i Oslo, Geografisk institutt. Naturgeografisk seksjon. Notat, 65 pp. (In Norwegian - on geologically valuable areas).

#### 6.1.2 - Additional reports and documents

i. taxonomic lists of plant and animal species occurring in the site (see section 4.3)

ii. a detailed Ecological Character Description (ECD) (in a national format)

iii. a description of the site in a national or regional wetland inventory

iv. relevant Article 3.2 reports

v. site management plan

<no file available>

vi. other published literature

#### 6.1.3 - Photograph(s) of the Site

Please provide at least one photograph of the site



A Nice day in the early summer in Kvisleflået ( Jor



Moosedung Moss ( Jo



Dry pine With Wolf Lichen



Bleketjønna ( Jon Bekken, 22-07-2017 )



Pair of Wooper Swans ( Jon Bekken, 22-06-2006 )

### 6.1.4 - Designation letter and related data

#### **Designation letter**

<1 file(s) uploaded>

Date of Designation 2002-08-06